BALCONES CANYONLANDS PRESERVE

LAND MANAGEMENT PLAN

TIER II-A



October 2015

Austin • Texas

Balcones Canyonlands Conservation Plan (BCCP) and the

Balcones Canyonlands Preserve (BCP)

The **Balcones Canyonlands Conservation Plan** (**BCCP**) is a federal Endangered Species Act (ESA) incidental "take" permit for 30 years issued to Travis County and the City of Austin on May 2, 1996 by the US Fish and Wildlife Service (USFWS). Incidental take is the loss of federally listed species or their habitats in the course of (or "incidental to") otherwise legal actions, like development. Such permitting is authorized under ESA Section 10(A)(1)(b), so sometimes the BCCP is called a "10A Permit."

A collection of documents guides BCCP implementation: our Endangered Species Act Permit No. TE 788841-2, the BCCP Final Environmental Impact Statement and Habitat Conservation Plan, the Travis County – City of Austin Interlocal Agreement – Shared Vision, Permit Area and Fee Zone Maps and **tiered Land Management Plans**.

These documents together provide the permit conditions, mitigation requirements, land acquisition areas (also known as the **Balcones Canyonlands Preserve** or **BCP**), management guidelines, and mechanisms by which the City and County can cover the impact of endangered species habitat loss in western Travis County and expedite development projects within the Permit Area.

The Land Management Plans are tiered:

Tier I	Overview of the Preserve and Partner Responsibilities
--------	---

- Tier II A BCP Land Management Guidelines (Specific Best Practices)
- Tier II B BCCP Administration
- Tier II C BCP Macrosite Requirements
- Tier III BCP individual tract plans

This plan outlines best practices for Karst Management, Tier II A-9.

TABLE OF CONTENTS

1.0 PURPOSE	1
2.0 BACKGROUND	1
2.1 COVERED SPECIES	1
2.2 FEATURES AND RELATIONSHIP TO BCCP SPECIES	3
3.0 NEW KARST INFORMATION RELATED TO THE BCCP	23
4.0 ADDITIONAL VALUES FOR CAVE AND KARST ECOSYSTEM	S 24
5.0 THREATS	25
6.0 MANAGEMENT PROGRAM	28
6.1 KARST MANAGEMENT GOALS	28
6.2 CONSERVATION ACTIONS	29
6.2.1 Vegetation Management	29
6.2.2 Animal Management	
6.2.3 Cave Gating and Fencing in BCP Caves and Bat Manager	nent32
6.2.4 Physical Management	34
6.2.5 Access Guidelines	35
6.2.6 Public Education and Outreach	36
6.3 MANAGEMENT COORDINATION	37
7.0 MONITORING / RESEARCH	38
7.1 NEWLY DISCOVERED KARST FEATURES IN THE PERMIT	AREA41
7.2 RESEARCH NEEDS	42
8.0 LITERATURE CITED	45

LIST OF TABLES

Table 1. Federally Listed Karst Species Covered by the BCCP1
Table 2. Karst Species of Concern Covered by the BCCP
Table 3. 62 BCCP Karst Features: Current Ownership Status
Table 4. Endangered Karst Invertebrate Locations within BCCP caves of Travis County, Texas
Table 5. Non-BCCP listed Caves/Karst Features with Listed Invertebrates Protected on BCP 14
Table 6. Karst Invertebrate SOC within BCCP Caves, Travis County, Texas 16
Table 7. Non-BCCP Caves/Karst Features with Karst SOC Protected on BCP21

LIST OF FIGURES

Figure 1. BCCP Cave Locations	5
Figure 2. Karst Clusters	6

LIST OF APPENDICES

Appendix A
BCCP Cave Substitution Policy

1.0 PURPOSE

This document outlines the policies and strategies for BCP cave and karst management; individual feature management specifics are outlined in the Tier III Land Management chapter for each BCP Unit or tract.

2.0 BACKGROUND

The regional ESA Section 10(a)(I)(B) permit (TE 788841-2), also known as the Balcones Canyonlands Conservation Plan (BCCP), requires the creation of the Balcones Canyonlands Preserve (BCP), protection of 62 karst features, and a high standard of protection, stewardship and adaptive management to secure habitat in perpetuity and protect populations of eight endangered species (ES) and 27 species of concern (SOC). The City of Austin and Travis County (Permit Holders), Lower Colorado River Authority (LCRA) and the City of Sunset Valley (Managing Partners), and other cooperating entities (e.g. private landowners, Travis Audubon, the City of Lakeway, Texas Cave Management Association, The Nature Conservancy of Texas) own and manage BCP species, habitats, and ecosystems.

2.1 COVERED SPECIES

Six endangered karst invertebrate species (**Table 1**) and 25 karst SOC are covered by the BCCP (**Table 2**). If these 25 SOC become federally listed as threatened or endangered, no additional mitigation by the Permit Holders would be required if all of the karst protection outlined in the BCCP is fully implemented.

Common Name	Scientific Name
Tooth Cave pseudoscorpion	Tartarocreagris texana
Tooth Cave spider	Tayshaneta myopica ¹
Tooth Cave ground beetle	Rhadine persephone
Kretschmarr Cave mold beetle	Texamaurops reddelli
Bee Creek Cave harvestman	Texella reddelli
Bone Cave harvestman	Texella reyesi

Footnotes follow table

¹ *Tayshaneta myopica* is listed in the regional permit as *Neoleptoneta myopica*, but a 2012 study revised the genus *Neoleptoneta*, thus identifying this species in the genus *Tayshaneta* (Campbell et al. 2012).

Common Name	Scientific Name
Flatworm	Sphalloplana mohri
Ostracod	Candona sp. nr. stagnalis
Isopod	Caecidotea reddelli
Isopod	Trichoniscinae N. S.
Isopod	Miktoniscus N. S.
Spider	Cicurina bandida
Spider	Cicurina cueva
Spider	Cicurina ellioti
Spider	Cicurina reddelli
Spider	Cicurina reyesi
Spider	Cicurina travisae
Spider	Cicurina wartoni
Spider	Tayshaneta concinna ¹
Spider	Tayshaneta devia ¹
Spider	Eidmannella reclusa
Pseudoscorpion	Aphrastochthonius N. S.
Pseudoscorpion	Tartarocreagris comanche ²
Pseudoscorpion	Tartarocreagris reddelli
Pseudoscorpion	Tartarocreagris intermedia
Pseudoscorpion	Tartarocreagris N. S. 3
Harvestman	Texella spinoperca
Millipede	Speodesmus N. S
Ground Beetle	Rhadine s. subterranea
Ground Beetle	Rhadine s. mitchelli
Ground Beetle	Rhadine austinica

Table 2. Karst Species of Concern Covered by the BCCP

¹ *Tayshaneta myopica* is listed in the regional permit as *Neoleptoneta myopica*, but a 2012 study revised the genus *Neoleptoneta*, thus identifying this species in the genus *Tayshaneta* (Campbell et al. 2012).

Footnotes follow table

² *Tartarocreagris comanche* is improperly listed in the regional permit as the New Comanche Trail Cave harvestman.

Species descriptions for endangered karst species known to occur in Travis County can be found in the Biological Advisory Team (BAT) report (1990), Recovery Plan for Endangered Karst Invertebrates in Travis and Williamson Counties, Texas (USFWS 1994), and USFWS 5-year reviews (USFWS 2008, 2009a, 2009b, 2009c).

2.2 FEATURES AND RELATIONSHIP TO BCCP SPECIES

Western Travis County is characterized as a strongly dissected limestone outcrop tableland, bordered abruptly on the east by the Balcones fault zone or Balcones Escarpment (Amos and Gehlbach 1988). In addition to surface habitat, the underlying karstic limestone, highly fractured and full of solution cavities, provides diverse subterranean habitats for specially adapted invertebrate and vertebrate species. The cave environment of central Texas, including that within the permit area, has been recognized to support one of the most important cave faunas in the world (Elliott and Reddell 1989).


The regional permit seeks to prevent the loss of caves known to contain federally endangered species covered by the Permit and includes protection for significant features, karst clusters, and preserves (**Figure 1: BCCP cave locations**). The regional permit, when fully implemented, will protect 35 of the 39 endangered species caves in Travis County that were known when the permit was issued in 1996. In addition, under the permit, 27 caves are proposed to be protected that support SOC for a total of 62 karst features to be protected under the BCCP. These SOC caves are recommended for protection because they support rare invertebrate species and are also important recharge features. These karst features provide water to be recharged to the Edwards Aquifer and help to protect the water quality of the Austin area. **Table 3** depicts **62 BCCP Karst Features - Current Ownership**.

Three cave clusters (**Figure 2: Karst Clusters**) have been identified within the BCCP permit area and also immediately outside the permit area to the northeast: the Four Points Cluster (includes the area northwest and northeast of the FM

2222/RM 620 intersection), the McNeil Cluster, and the Northwood Cluster. The Northwood and McNeil clusters occur in close proximity to each other in the vicinity of Walnut Creek near Howard Lane and McNeil Drive in North Austin. Twenty-seven of the 62 karst features (62 = 60 caves, one spring, and one mine) covered by this Karst Management Plan are privately owned (**Table 3**). BCP Partners work with willing non-profit groups, private landowners and other interested parties to protect these privately owned listed karst features.

Known species distribution in the BCCP-listed caves and caves not listed on the Permit, but protected in the BCP, are depicted in Table 4 through Table 7.

The environmental integrity of all 62 karst features is proposed to be protected through acquisition and management, or implementation of а management/conservation agreement with entities that influence the hydrogeological area needed to protect the feature (USFWS 1996a). Management in karst preserves includes maintenance of native vegetation, redimported fire ant (RIFA) control, control of disturbance by humans, and protection of water quality and nutrient input.








Figure 2. Karst Clusters

Table 3. 62 BCCP Karst Features: Current Ownership Status Key and footnotes follow table

Cave Name	ES or SOC	Current tract/owner In BCP or Private	Cave Cluster
Adobe Springs Cave	SOC	BCP Lehmann/TNC	
Airman's Cave	SOC	BCP Barton Creek/COA	
Amber Cave	ES	BCP Jollyville/TC	Four Points (West)
Armadillo Ranch Sink	SOC	Private	
Arrow Cave	SOC	BCP Slaughter Creek Park./COA	
Bandit Cave	ES	Private	
Beard Ranch Cave (Featherman's Cave)	ES	BCP Ivanhoe/COA	
Bee Creek Cave	ES	Private	
Blowing Sink Cave	SOC	BCP COA	
Broken Arrow Cave	ES	BCP Lime Creek Preserve/COA	
Buda Boulder Spring	SOC	BCP Shoal Creek Greenbelt/COA	
Cave X	SOC	Private/COA Protection Agreement	
Cave Y ¹	SOC	BCP Barton Creek Greenbelt/COA	
Ceiling Slot Cave	SOC	Private	
Cold Cave	ES	Private	Northwood
Cotterell Cave	ES	BCP Stillhouse Hollow Preserve/COA	
Disbelievers Cave	ES	BCP Private 10(a)	Four Points (East)
District Park Cave	SOC	BCP Dick Nickols Park/COA	
Eluvial Cave	ES	BCP Private 10(a)	Four Points (East)
Flint Ridge Cave	SOC	Prop 2 Tabor Tract /COA	
Fossil Cave	ES	BCP Schroeter Park/COA	
Fossil Garden Cave	ES	Private	McNeil
Gallifer Cave ES BCP Jollyville/TC		BCP Jollyville/TC	Four Points (West)

Cave Name	ES or SOC	Current tract/owner In BCP or Private	Cave Cluster
Get Down Cave	SOC	Private/COA Protection Agreement	
Goat Cave	SOC	BCP Goat Cave Karst Preserve/COA	
Hole-in-the-Road Cave	ES	Private	Northwood
Ireland's Cave	SOC	BCP Ireland's/ TC	
Jack's Joint	SOC	Private	
Japygid Cave	ES	BCP Private 10(a)	Four Points (East)
Jest John Cave	ES	BCP Forest Ridge/COA	
Jester Estates Cave	ES	BCP Forest Ridge/COA	
Jollyville Plateau Cave	ES	BCP Private 10(a)	Four Points (East)
Kretschmarr Cave	ES	BCP Jollyville/TC	Four Points (West)
Kretschmarr Double Pit	ES	BCP Jollyville/TC	Four Points (West)
Lamm Cave	ES	BCP Private Section 7	
Little Bee Creek Cave	ES	BCP Ullrich WTP/COA	
Lost Gold Cave	SOC	Private	
Lost Oasis Cave	SOC	Private TCMA	
M.W.A. Cave	ES	BCP Private 10(a)	Four Points (East)
Maple Run Cave	SOC	BCP Goat Cave Karst Preserve/COA	
McDonald Cave	ES	BCP Jollyville/TC	
McNeil Bat Cave	ES	Private	McNeil
Midnight Cave	SOC	BCP Slaughter Creek Park/COA	
Moss Pit	SOC	Private	
New Comanche Trail Cave	ES	BCP Lake Travis/TC	
No Rent Cave	ES	Private	McNeil
North Root Cave	ES	BCP Jollyville/TC	Four Points (West)
Pennie's Cave	SOC	Private	

Cave Name	ES or SOC	Current tract/owner In BCP or Private	Cave Cluster
Pickle Pit	SOC	BCP Private Section 7	
Pipeline Cave	SOC	Private	
Rolling Rock Cave	ES	BCP Lime Creek Preserve/COA	
Root Cave	ES	BCP Jollyville/TC	Four Points (West)
Slaughter Creek Cave	SOC	BCP Slaughter Creek Park/COA	
Spanish Wells Cave	SOC	BCP Kotrla/TC	
Spider Cave	ES	BCP Park West/COA	
Stark's North Mine ²	ES	BCP Stark's/ TC	
Stovepipe Cave	ES	BCP Canyon Creek/ COA	
Talus SpringsN/ACave3		Private/ 10(a) permit	
Tardus Hole	ES	BCP Jollyville/TC	Four Points (West)
Tooth Cave	ES	BCP Jollyville/TC	Four Points (West)
Weldon Cave	ES	Private	McNeil
Whirlpool Cave SOC		Private TCMA	

Key and Footnotes

- ES = Endangered (federally listed) Species
- SOC = Species of Concern

¹Cave Y was considered an ES cave *(Texella reddelli)* in the 1996 BCCP Permit, but has since been determined not to contain *Texella reddelli* (Reddell 2004).

²Stark's North Mine was listed as a SOC cave in the 1996 BCCP Permit, but has since been determined to contain *Texella reddelli* (USFWS 2009c).

³ Talus Springs Cave has never been known to contain ES or SOC (Elliot 1997).

Table 4. Endangered Karst Invertebrate Locations within BCCP caves of Travis County, Texas

This table, originally in the BCCP 1996 documents, has been revised to show new species' location information. Key and footnotes follow table

Cave Name	Current Preserve Status	Karst Fauna Region	Tooth Cave Pseudoscorpion Tartarocreagris texana	Tooth Cave Spider Tayshaneta myopica	Tooth Cave Ground Beetle <i>Rhadine</i> <i>persephone</i>	Kretschmarr Cave Mold Beetle <i>Texamaurops</i> <i>reddelli</i>	Bee Creek Cave Harvestman <i>Texella</i> reddelli	Bone Cave Harvestman <i>Texella reyesi</i>
Amber Cave	BCP Jollyville TC	Jollyville Plateau	X 1996		X 2010 (Reddell)	X 1996		
Bandit Cave	Private	Rolling- wood					P 1996 X 2009	
Beard Ranch Cave	BCP Ivanhoe COA	Jollyville Plateau						X 1996
Bee Creek Cave	Private	Rolling- wood					X 1996	
Broken Arrow Cave	BCP Lime Creek Preserve COA	Cedar Park			X 1996			
Cold Cave	Private	McNeil Round Rock						X 1996
Cotterell Cave	BCP Spicewood Springs Park/COA	Central Austin						X 1996
Disbelievers Cave	BCP Private 10(a)	Jollyville			X 1996			
Eluvial Cave	BCP Private 10(a)	Jollyville						X 1996
Fossil Cave	BCP Schroeter Park/COA	McNeil Round Rock						X 1996

Cave Name	Current Preserve Status	Karst Fauna Region	Tooth Cave Pseudoscorpion Tartarocreagris texana	Tooth Cave Spider Tayshaneta myopica	Tooth Cave Ground Beetle <i>Rhadine</i> <i>persephone</i>	Kretschmarr Cave Mold Beetle <i>Texamaurops</i> <i>reddelli</i>	Bee Creek Cave Harvestman <i>Texella</i> <i>reddelli</i>	Bone Cave Harvestman <i>Texella reyesi</i>
Fossil Garden Cave	Private	McNeil Round Rock						X 1996
Gallifer Cave	BCP Jollyville/TC	Jollyville Plateau		P 1996 X 2010 (Ledford)	P 1996 X 2005	X 2009 (Chandler)		X 1996
Hole-in-the- Road Cave	Private	McNeil Round Rock						X 1996
Japygid Cave	BCP Private 10(a)	Jollyville			X 1996	P 1996 X 2005		
Jest John Cave	BCP Forest Ridge/COA	Jollyville Plateau					X 1996	
Jester Estates Cave	BCP Forest Ridge/COA	Jollyville Plateau	X 2008 (Cokendolpher)	X 2010 (Ledford)			X 1996	
Jollyville Plateau Cave	BCP Private 10(a)	Jollyville			X 1996			X 1996
Kretschmarr Cave	BCP Jollyville/TC	Jollyville Plateau			X 1996	X 1996		
Kretschmarr Double Pit	BCP Jollyville/TC	Jollyville Plateau	P 1996 X 2005		P 1996 X 2005		P 1996	
Lamm Cave	Private Section 7	Jollyville Plateau			X 1996			
Little Bee Creek Cave	BCP Ullrich WTP/COA	Rolling- wood					X 1996	
McDonald Cave	BCP Jollyville/TC	Jollyville Plateau						X 1996
McNeil Bat Cave	Private	McNeil Round Rock		X 2010 (Ledford)				X 1996

Cave Name	Current Preserve Status	Karst Fauna Region	Tooth Cave Pseudoscorpion <i>Tartarocreagris</i> <i>texana</i>	Tooth Cave Spider Tayshaneta myopica	Tooth Cave Ground Beetle <i>Rhadine</i> <i>persephone</i>	Kretschmarr Cave Mold Beetle <i>Texamaurops</i> <i>reddelli</i>	Bee Creek Cave Harvestman <i>Texella</i> <i>reddelli</i>	Bone Cave Harvestman <i>Texella reyesi</i>
M.W.A. Cave	BCP Private 10(a)	Jollyville	P 1996 X 2005		X 1996	P 1996 X 2005		X 1996
New Comanche Trail Cave	BCP Lake Travis/TC	Jollyville Plateau		X 1996				X 1996
No Rent Cave	Private	McNeil Round Rock						X 1996
North Root Cave	BCP Jollyville/TC	Jollyville Plateau			X 1996			
Rolling Rock Cave	BCP Lime Creek Preserve COA, Sec.10(a)	Cedar Park			X 1996			
Root Cave	BCP Jollyville/TC	Jollyville Plateau		X 2010 (Ledford)	X 1996			X 1996
Spider Cave	BCP Park West/COA	Jollyville Plateau			X 2004 (Reddell)		X 2004 (Reddell)	
Stark's North Mine Cave	BCP Stark's/TC						X 2009 (USFWS)	
Stovepipe Cave	BCP Canyon Creek/ COA	Jollyville Plateau			X 1996	X 1996		P 1996 X 2009 (USFWS)
Tardus Hole	BCP Jollyville/TC	Jollyville Plateau			X 1996	X 2009 (Chandler)		
Tooth Cave	BCP Jollyville/TC	Jollyville Plateau	X 1996	X 1996	X 1996	X 1996		X 1996
Weldon Cave	Private	McNeil Round Rock						X 1996

Sources: BCCP Permit 1996, Elliott 1992, USFWS 1994, Reddell 2004, 2005, 2010, HNTB 2005, USFWS 2009a, 2009b, 2009c, Ledford 2010. Key and Footnotes

- X 1996 = confirmed occurrence based on collected specimen, the designation in the 1996 BCCP permit
- P 1996 = probable occurrence based on observation but not confirmed with collected specimen, the designation in 1996 BCCP permit
- X 2005 = was listed as confirmed in the HNTB summary of James Reddell's data, 2005 report for USFWS
- X 2008 = Cokendolpher (pers com 2008) confirmed that Jester Estates Cave is a new site for Tartarocreagris texana

X 2009 = USFWS - according to the 2009 5 year review on *Texella reyesi* the report lists *T. reyesi* as confirmed for Stovepipe Cave; *Texella reddelli* 5-year review confirms *T. reddelli* for Bandit Cave and Stark's North Mine (USFWS 2009c).

- X 2009 (Chandler) = confirmed by D. Chandler, as reported in USFWS 5-year review (2009b).
- X 2010 (Ledford) = confirmed by J. Ledford (pers com 2010)
- X 2010 (Reddell) = confirmed by J. Reddell (pers com 2010)

Table 5. Non-BCCP listed Caves/Karst Features with Listed Invertebrates Protected on BCP

Cave Name	Current Preserve Status	Karst Fauna Region	Tooth Cave Pseudoscorpion Tartarocreagris texana	Tooth Cave Spider Tayshaneta myopica	Tooth Cave Ground Beetle <i>Rhadine</i> <i>persephone</i>	Kretschmarr Cave Mold Beetle <i>Texamaurops</i> <i>reddelli</i>	Bee Creek Cave Harvestman <i>Texella</i> <i>reddelli</i>	Bone Cave Harvestman <i>Texella</i> <i>reyesi</i>
Cortana Cave	COA	Jollyville Plateau		X 2010				X 2008
Down Dip Sink	COA	Jollyville Plateau			X 2007a			
Garden Hoe Cave	COA	Jollyville Plateau			X 2007b			
Geode Cave	тс	Jollyville Plateau		X 2008	X 2008			X 2008
LU-11	тс	Jollyville Plateau		X 2008				
LU-12	тс	Jollyville Plateau						X 2008
IV-3	COA	Jollyville Plateau						X 2010
Little Black Hole	COA	Rollingwood					X 2009c	
Merkin Hole	COA	Jollyville Plateau					X 2010	
Pond Party Pit	COA	Jollyville Plateau						X 2010
RI-1	TC	Jollyville Plateau					X 2010	

Key follows table

Cave Name	Current Preserve Status	Karst Fauna Region	Tooth Cave Pseudoscorpion Tartarocreagris texana	Tooth Cave Spider <i>Tayshaneta</i> <i>myopica</i>	Tooth Cave Ground Beetle <i>Rhadine</i> <i>persephone</i>	Kretschmarr Cave Mold Beetle <i>Texamaurops</i> <i>reddelli</i>	Bee Creek Cave Harvestman <i>Texella</i> reddelli	Bone Cave Harvestman <i>Texella</i> <i>reyesi</i>
Tight Pit Cave	тс	Jollyville Plateau		X 2010				
Two Trunks Cave	TC	Jollyville Plateau			X 2008 (USFWS)			

Sources: USFWS 2008, 2009c, Zara Environmental 2007a, 2007b, 2008, and 2010.

<u>Key</u>

X = confirmed occurrence based on collected specimen.

Table 6. Karst Invertebrate SOC within BCCP Caves, Travis County, Texas^{1,2}

Key and footnotes follow table

Cave Name	Aphrastochithonius N.S.	Caecidotea reddelli	Candona sp. nr. stagnalis	Cicurina bandida ³	Cicurina travisae ⁴	Cicurina sp. ⁵	Eidmannella reclusa	Miktoniscus N.S.	Tayshaneta concinna	Tayshaneta devia	Rhadine austinica	Rhadine s. subterranea	Rhadine s. mitchelli	Speodesmus N.S.	Sphalloplana mohri	Tartarocreagris comanche	Tartarocreagris intermedia	Tartarocreagris N.S. 3	Texella spinoperca	Trichoniscinae N.S.
Adobe Springs Cave																				
Airmen's Cave				Х							Х						Х		Х	
Amber Cave					Х								Х							
Armadillo Ranch Sink		Х																		
Arrow Cave				Х							Х									
Bandit Cave				Х							Х									Х
Beard Ranch Cave					Х															
Bee Creek Cave				Х							Х									
Blowing Sink Cave				Х							Х									
Broken Arrow Cave																				
Buda Boulder Spring		Х																		
Cave X		Х	Х	Х				Х			Х			Х						
Cave Y				Х							Х									

Cave Name	Aphrastochithonius N.S.	Caecidotea reddelli	Candona sp. nr. stagnalis	Cicurina bandida ³	Cicurina travisae ⁴	Cicurina sp. ⁵	Eidmannella reclusa	Miktoniscus N.S.	Tayshaneta concinna	Tayshaneta devia	Rhadine austinica	Rhadine s. subterranea	Rhadine s. mitchelli	Speodesmus N.S.	Sphalloplana mohri	Tartarocreagris comanche	Tartarocreagris intermedia	Tartarocreagris N.S. 3	Texella spinoperca	Trichoniscinae N.S.
Ceiling Slot Cave						Х														
Cold Cave						Х														
Cotterell Cave					Х							Х								
Disbelievers Cave						Х														
District Park Cave				Х							Х									
Eluvial Cave																				
Flint Ridge Cave				Х							Х									
Fossil Cave												Х								
Fossil Garden Cave						Х						Х								
Gallifer Cave					Х															
Get Down Cave				Х							Х									
Goat Cave				Х										Х						
Hole-in-the- Road Cave						х														
Ireland's Cave				Х							Х									
Jack's Joint		Х				Х														
Japygid Cave																				

Cave Name	Aphrastochithonius N.S.	Caecidotea reddelli	Candona sp. nr. stagnalis	Cicurina bandida ³	Cicurina travisae ⁴	Cicurina sp. ⁵	Eidmannella reclusa	Miktoniscus N.S.	Tayshaneta concinna	Tayshaneta devia	Rhadine austinica	Rhadine s. subterranea	Rhadine s. mitchelli	Speodesmus N.S.	Sphalloplana mohri	Tartarocreagris comanche	Tartarocreagris intermedia	Tartarocreagris N.S. 3	Texella spinoperca	Trichoniscinae N.S.
Jest John Cave					Х															
Jester Estates Cave					Х															
Jollyville Plateau Cave																				
Kretschmarr Cave					Х								Х							
Kretschmarr Double Pit					Х															
Lamm Cave																				
Little Bee Creek Cave				Х							Х									
Lost Gold Cave				х					Х		Х									
Lost Oasis Cave				х							Х									
M.W.A. Cave																		Х		
Maple Run Cave				Х							Х									
McDonald Cave					Х					Х										
McNeil Bat Cave												Х								
Midnight Cave				Х							Х									

Cave Name	Aphrastochithonius N.S.	Caecidotea reddelli	Candona sp. nr. stagnalis	Cicurina bandida ³	Cicurina travisae ⁴	Cicurina sp. ⁵	Eidmannella reclusa	Miktoniscus N.S.	Tayshaneta concinna	Tayshaneta devia	Rhadine austinica	Rhadine s. subterranea	Rhadine s. mitchelli	Speodesmus N.S.	Sphalloplana mohri	Tartarocreagris comanche	Tartarocreagris intermedia	Tartarocreagris N.S. 3	Texella spinoperca	Trichoniscinae N.S.
Moss Pit																				
New Comanche Trail Cave						x	х									х				
No Rent Cave												Х								
North Root Cave					х															
Pennie's Cave											Х			Х						
Pickle Pit					Х															
Pipeline Cave														Х						
Rolling Rock Cave						Х								Х						
Root Cave					Х															
Slaughter Creek Cave				Х																
Spanish Wells Cave		х													Х					
Spider Cave					Х		Х													
Stark's North Mine									Х											
Stovepipe Cave	Х				Х		Х													
Talus Springs Cave																				

Cave Name	Aphrastochithonius N.S.	Caecidotea reddelli	Candona sp. nr. stagnalis	Cicurina bandida ³	Cicurina travisae ⁴	Cicurina sp. ⁵	Eidmannella reclusa	Miktoniscus N.S.	Tayshaneta concinna	Tayshaneta devia	Rhadine austinica	Rhadine s. subterranea	Rhadine s. mitchelli	Speodesmus N.S.	Sphalloplana mohri	Tartarocreagris comanche	Tartarocreagris intermedia	Tartarocreagris N.S. 3	Texella spinoperca	Trichoniscinae N.S.
Tardus Hole																				
Tooth Cave					Х		Х						Х							
Weldon Cave	Х											Х								
Whirlpool Cave				Х							Х									

Sources: Elliot 1997, Paquin and Hedin 2005, Paquin et al. 2008, TMM 2007, Zara Environmental 2008, 2010, Hedin 2015.

Key and Footnotes

X = confirmed location based on collected specimen.

¹ *Cicurina ellioti* listed as an SOC in the regional permit is not included in this table because this species has now been synonymized with Cicurina buwata, a non-SOC (Cokendolpher 2004).

² *Tartarocreagris reddelli* listed as a SOC in the regional permit is not included in this table because this species has now been synonymized with *Tartarocreagris infernalis*, a non-SOC (Muchmore 2001).

³ Occurrences of *Cicurina bandida* include localities formerly listed as *Cicurina cueva* and *Cicurina reyesi*, which have been formally grouped together into the single species *C. bandida* (Paquin et al. 2008).

⁴ Occurrences of *Cicurina travisae* include localities formerly listed as *Cicurina reddelli* and *Cicurina wartoni*, which have been formally grouped together into the single species *C. travisae* (Hedin 2015).

⁵Localities of possible SOCs; blind *Cicurina* specimens not yet confirmed to species level.

Table 7. Non-BCCP Caves/Karst Features with Karst SOC Protected on BCP^{1,2}

Key and footnotes follow table

Cave Name	BCP Owner	Aphrastochithonius N.S.	Caecidotea reddelli	Candona sp. nr. stagnalis	Cicurina bandida ³	Cicurina travisae ⁴	Cicurina sp. ^s	Eidmannella reclusa	Miktoniscus N.S.	Tayshaneta concinna	Tayshaneta devia	Rhadine austinica	Rhadine s. subterranea	Rhadine s. mitchelli	Speodesmus N.S.	Sphalloplana mohri	Tartarocreagris comanche	Tartarocreagris intermedia	Tartarocreagris N.S. 3	Texella spinoperca	Trichoniscinae N.S.
Brewpot Cave	тс										Х										
Cortana Cave	COA						Х														
Down Dip Cave	COA						Х														
Geode Cave	тс					Х		Х													
IV-3	COA					Х															
LU-29	тс						Х														
Pond Party Pit	COA						Х														
RI-1	тс						Х														
RI-3	тс						Х														
Siebert Sink	COA				х					Х										х	
Two Trunks Cave	тс					х															

Sources: Bayless pers com 2013, Paquin and Hedin 2005, Sanders pers com 2013, TMM 2007, Zara Environmental 2008, 2010, Hedin 2015.

BCP Land Management Plan

Key and Footnotes

X =confirmed location based on collected specimen.

¹ *Cicurina ellioti* listed as an SOC in the regional permit is not included in this table because this species has now been synonymized with Cicurina buwata, a non-SOC (Cokendolpher 2004).

² *Tartarocreagris reddelli* listed as a SOC in the regional permit is not included in this table because this species has now been synonymized with *Tartarocreagris infernalis*, a non-SOC (Muchmore 2001).

³ Occurrences of *Cicurina bandida* include localities formerly listed as *Cicurina cueva* and *Cicurina reyesi*, which have been formally grouped together into the single species *C. bandida* (Paquin et al. 2008).

⁴ Occurrences of *Cicurina travisae* include localities formerly listed as *Cicurina reddelli* and *Cicurina wartoni*, which have been formally grouped together into the single species *C. travisae* (Hedin 2015).

⁵Localities of possible SOCs; blind *Cicurina* specimens not yet confirmed to species level.

Tier II A Chapter 9 Karst Management

3.0 NEW KARST INFORMATION RELATED TO THE BCCP

For 18 years, "The Caves of the Balcones Canyonlands Conservation Plan, Travis County, Texas" (Elliot 1997) has been the primary reference guide for endangered species and SOC location information. Recently, however, the USFWS released 5-year reviews for the six endangered karst species listed on the BCCP permit (USFWS 2008, 2009a, 2009b), which included documentation of new localities for these species. More recent survey work by Zara Environmental, Inc. (2007a, 2007b, 2008, 2010) has also added new location information for four of the endangered karst species and several SOCs listed on the BCCP permit. A complete list of known endangered karst invertebrate locations for the BCCP-listed karst features is summarized in Table 2; known SOC localities within BCCP-listed karst features are summarized in Table 4. Location information for endangered karst species and SOCs found in BCP caves that were not listed in the BCCP are summarized in Table 3 (ES localities) and Table 5 (SOC localities). Though not listed on the permit, these caves and any other BCP caves containing ES or SOC found in the future will be protected in the same manner as those listed on the permit.

Joel Ledford (University of California, Berkeley) conducted a revision of the Family Leptonetidae with particular emphasis on the taxonomy and relationships within the subfamily Archoleptonetinae. This study found new locations for the endangered *Neoleptoneta myopica* and newly described species within the Austin area. Ledford also proposed to change the genus *Neoleptoneta* to *Tayshaneta* (Campbell et al. 2012). USFWS adopted this change in 2015 (Watson pers com 2015).

Marshal Hedin (San Diego State University) conducted a study for USFWS using DNA sequence data to rigorously test the species status of *Cicurina wartoni*, a BCCP-listed SOC known only from Pickle Pit Cave. Hedin's study used specimens collected by BCP staff from multiple caves in northern Travis and southern Williamson counties to determine if *Cicurina* specimens from Pickle Pit were genetically distinguishable from other nearby sites containing *Cicurina*

spiders previously identified as *C. buwata* (formerly known as *C. ellioti*), *C. reddelli*, and *C. travisae*. Results of genetic analyses indicate that there are only two distinct species complexes in the study area: *C. buwata* in the northern range and *C. travisae* in the southern range. Based on these findings, *C. reddelli*, *C. wartoni*, and *C. travisae* should now be treated as a single species: *C. travisae*. Thus, confirmed localities previously identified as *C. reddelli* and *C. wartoni* are now considered as localities for *C. travisae* (Hedin 2015). Following the completion of Hedin's study, USFWS completed a status review of *Cicurina wartoni* and concluded that this species does not warrant protection under the Endangered Species Act (ESA) (USFWS 2014a).

4.0 ADDITIONAL VALUES FOR CAVE AND KARST ECOSYSTEMS

Beyond protecting the entrances of caves that are localities for endangered karst invertebrates and SOCs, USFWS Karst Preserve Design Recommendations (2012) also describe the importance of protecting the surface environment surrounding caves. One component of this protection involves preserving adequate habitat for trogloxenes such as cave crickets, bats, and mammals (USFWS 2012). Cave crickets are considered a keystone species for cave ecosystems, providing vital nutrients into an otherwise nutrient poor environment (Taylor et al. 2005). Bats and mammals such as raccoons are also important biotic components of karst ecosystems, supplying nutrient input in the forms of guano and scat which benefits resident karst invertebrates (USFWS 2011e). Providing adequate protection of surface plant and animal communities in cave preserves benefits these trogloxenes, and also protects other sources of nutrient input in the form of roots, leaf-litter, and woody debris, thereby creating a higher probability of long-term survival for protected karst invertebrates (USFWS 2012).

Another component of protecting the surface environment around caves involves maintaining high quality and adequate quantity of water to the cave ecosystem, achieved through protection of a cave's surface and sub-surface drainage basins (USFWS 2012). Well protected drainage basins provide necessary moisture and

stable temperatures in cave habitats, and ensure these ecosystems are free from contaminants (USFWS 2012).

5.0 THREATS

One of the main threats to the listed karst species is loss of habitat due to urban development activities. These species occur in an area that is undergoing continued urban expansion at a rapid rate and few caves are protected. Most of the species' localities occur adjacent to or near developed areas, or in areas that are proposed for development (USFWS 1996a).

The most significant effects of urban development on karst habitat are:

- filling of cave entrances or greatly reduced infiltration due to impervious cover. This blockage decreases the total energy entering the cave through the entrance (Russell pers com 1998) and reduces the moisture input necessary to maintain high humidity in the cave.
- inadequate setbacks for cave cricket foraging areas. Vital nutrient input provided by cave crickets could be lost if efforts are not made to protect their entire foraging range (105 meter radius around the cave footprint) (Taylor et al. 2005).
- pollutants from urban run-off, such as pesticides, which can contaminate caves and possibly harm or kill karst species or species that provide organic matter. Urban run-off can also alter the natural flow of nutrients through the cave system by replacing water flow and animal energy inputs with potentially contaminated seepage from yards and parking lots. If the surface and sub-surface drainage basins are not adequately protected, contamination of this nature can be expected.

Other threats to caves related to urban development include alteration of surface plant and animal communities, increased human visitation, vandalism, dumping, habitat fragmentation, and poorly designed cave gates (USFWS 2011a, 2011b). Land use changes can also affect the abundance and spatial arrangement of other organisms in the surface and sub-surface biotic community known to be beneficial to karst invertebrates (USFWS 2011a). Neglect of caves is also a threat since caves that are not visited or monitored may deteriorate due to inattention to new developments in cave areas; also, cave locations may be lost. Activities at several limestone quarries in northwestern Travis County may also threaten to destroy surrounding karst habitat. (BAT 1990, USFWS 1994).

Twenty percent of the known caves in Travis County have been covered or destroyed in the 20 years prior to the establishment of the BCP as a result of land use practices and development. This rate of loss is expected to continue (USFWS 1996a).

Recent scientific evidence of climate change demonstrates increases in average air temperatures in the last 50 years, coupled with an increase in heat waves and heavy precipitation events (IPCC 2007). These trends are projected to continue and increase in the next century with the southwestern U.S. being the most impacted of the continental U.S. (IPCC 2007). Karst invertebrates may be affected by the effects of climate change, due to their dependence on stable temperatures and humidity (USFWS 2011a). Climate change may impact karst species directly from increased in-cave temperatures and indirectly through changes in the vegetation and surface environment, which could affect food resource availability (USFWS 2011a). The caves of the Jollyville Plateau may be especially vulnerable to global warming due to the fact that they are shallow (generally 20 to 30 feet in depth). Rainfall regime changes and more extreme rain events may also impact the cave environments by flooding, filling in with debris, or adversely affecting nutrient inputs (USFWS 2011a).

Red imported fire ants (*Solenopsis invicta*) (RIFA) threaten the karst community directly by preying on karst invertebrates, but could also indirectly threaten them by reducing the amount of organic nutrients brought in by trogloxene species (species that live in the cave during the day and venture out at night foraging for food). Most notable trogloxenes are cave crickets and mammals such as raccoons. If the cave is overrun by RIFA, trogloxenes may disappear. RIFA will eat cave cricket eggs, nymphs and adults as well as forcing out mammals, greatly reducing the availability of organic material entering the cave. RIFA are

most abundant in disturbed areas (USFWS 2011b). Current estimates indicate that most of the 62 caves have at least some RIFA activity (Sanders pers com 2013; Bayless pers com 2013). See Tier II-A, Chapter X for additional information on RIFA.

Tawny crazy ants (*Nylanderia fulva*) are the latest invasive non-native species to threaten karst invertebrates. These newly arrived, non-native ants are a poorly understood species in the Austin area, making it difficult to project what long term impacts this species may have on karst ecosystems. In the Houston area this species has proven to be a major pest, and in areas of heavy infestation they have displaced RIFA (Meyers 2008). This species will likely have adverse effects on ant diversity as well as abundance and diversity of other arthropods in infestation areas (Meyers 2008). Since tawny crazy ants prefer wetter, more humid environments (Meyers 2008), areas around caves may be even more susceptible to invading colonies by providing preferred habitat characteristics. As of July , 2013 tawny crazy ants were confirmed at the entrance of Whirlpool Cave, and documented foraging as far as 100 ft inside the cave itself (Sanders pers com 2013; Bayless pers com 2013). TCAs were also documented infesting No Rent Cave in November 2014 (Sanders pers com 2015; Bayless pers com 2015).

Mammals bring in tremendous amounts of organic material into caves via their scat. Although the endangered karst fauna are very much dependent on these species to provide this material, the effects of large amounts of scat can also be detrimental when they attract non-cave adapted species (i.e. roaches) (Reddell pers com 2004).

White nose syndrome (WNS) is a newly observed disease responsible for unprecedented mortality of hibernating bats in the northeastern U.S., and since its discovery in 2007 has spread rapidly westward, posing a serious threat to hibernating bats throughout North America (USFWS 2011d). One species that commonly occurs in Travis County, the tri-colored bat (*Perimyotis subflavus*), has been shown to be susceptible to WNS (USFWS 2011d). In the 2013-2014

monitoring season (winter), WNS was detected and confirmed in Arkansas (USFWS 2014b). This occurrence demonstrates the potential for WNS to spread into the western U.S. in the near future (USFWS 2011d). Therefore, the threat of WNS to these important trogloxenes requires special attention of researchers accessing caves to be aware of potential transmission of this disease as well as appropriate decontamination procedures if WNS finds its way into central Texas caves (USFWS 2011d).

6.0 MANAGEMENT PROGRAM

6.1 KARST MANAGEMENT GOALS

The Recovery Plan for Endangered Karst Invertebrates in Travis County, Texas (USFWS 1994) outlines four major recovery actions: (1) research and information needs, (2) long-term protection for karst fauna areas, (3) monitoring, and (4) education. The BCCP's Habitat Conservation Plan/Environmental Impact Statement states that the BCCP should effectively implement these goals in order to assure that the implementation of the BCCP has no negative impact on the population viability of the endangered karst invertebrates (USFWS 1996a). Karst preserve design is the most important aspect for guarantying the long term survival of the species. Preserves that have adequate setbacks to ensure that the entire surface and subsurface drainage basins as well as the native plant and animal communities are protected will greatly enhance the long term success of the program (UFSWS 2012). The ultimate goal is wherever possible to have quality preserves that are self-sustaining, thus greatly reducing the need for intensive onsite management. A secondary management goal includes the protection of the BCP karst features to protect local water quality.

Currently protected karst habitat will be maintained and enhanced, and permit holders will attempt to protect or acquire additional BCCP caves for karst preserves. BCP partners will attempt to enter into formal management agreement(s) with the landowner(s) for all caves that are recommended for protection but have yet to be acquired or kept in private ownership as cave preserves. The management agreement(s) will detail the area to be managed for cave protection, what such management will entail, and who is responsible for the management. Efforts are needed to increase public awareness and sensitivity to the karst invertebrates and other endangered species.

6.2 CONSERVATION ACTIONS

The following is a summary of more detailed management information available from current literature, TC Natural Resources and Environmental Quality Division, COA - Austin Water Utility, and the USFWS. The following activities will be undertaken for caves owned or managed by BCP partner agencies for the 62 BCCP caves, as well as other BCP caves with ES or SOC.

If monitoring data shows that management methods are ineffective or can be improved, permit holders should practice "adaptive management"; in these cases the management plan will be revised and/or additional activities will be added. Such additions may include: fencing of additional areas around caves to control access, more intensive RIFA control, removal of non-native plant/animal species found to be detrimental to the karst ecosystem, or removal of additional species found to directly harm the species either directly (e.g. predators) or indirectly (e.g. species that prey on food base or increase the nutrient level (e.g. large amounts of raccoon scat attracting more aggressive surface species into the cave).

6.2.1 Vegetation Management

Ashe juniper-oak woodlands and other native vegetation will be protected within the preserve areas. Thick vegetation will be left to help protect caves by camouflaging their entrances. The size of the surface area needed to protect individual caves will be determined based on karst preserve design recommendations (USFWS 2012). Non-native vegetation in the critical area around a cave will be controlled to protect the cave ecosystem, preferably by mechanical control methods (USFWS 2011b). If chemical control methods to eliminate non-native plants around caves are absolutely necessary, herbicide treatments will be limited to cut-stump methods only (applying herbicide individually to freshly cut stumps or stems, which eliminates potential of drift and run-off); no foliar spray treatments will be used within the 105 m cave cricket foraging area of ES/SOC caves, or within the surface or subsurface drainage basin if run-off is potentially an issue.

When possible, the permit holders will work with nearby developers and landowners in the cave vicinity to encourage xeriscaped landscaping using native plants, which promotes less watering, fertilizers or pesticides, thereby minimizing groundwater contamination. Permit holders will also discourage the presence of non-native fauna such as feral hogs, which may damage native vegetation on cave surfaces.

6.2.2 Animal Management

RIFA should be controlled using USFWS approved methods (USFWS 2011b; see also Tier II-A, Chapter X). Surveys for RIFA mounds should be conducted at least twice per year. RIFA do not maintain their mounds during the summer, making them more difficult to see, but begin rebuilding them as soon as rains and cooler temperatures return (Vinson and Sorensen 1986). Because of this, at least one monitoring survey should be conducted in both spring and fall. These surveys should be conducted over the minimum cave cricket foraging area (within 80 m (262 ft) of cave entrances) and should be sufficient to yield actual RIFA mound densities, not merely indices of RIFA density. In addition, every routine maintenance inspection should include a search for RIFA mounds within 10 m (33 ft) of the cave entrance (USFWS 2011b). To avoid impacting the native ant population, the site must be surveyed for the presence of native ants prior to any RIFA treatment.

Control of RIFA should also be conducted at least twice per year if monitoring indicates their presence. RIFA may remain relatively inactive and deep within their mounds during long periods of drought or cold (Vinson and Sorensen 1986), making them more difficult to eradicate. Because of this, RIFA control should be conducted at least once in the spring and at least once in the fall. This control should be done shortly after the scheduled monitoring and not before so as not to artificially reduce the apparent RIFA density. An increase in the frequency of

RIFA control may be necessary based on (1) declines in cave cricket abundance or (2) an increase in the number of RIFA mounds within 80 m of the cave entrance (USFWS 2011b). Additionally, if RIFA mounds are observed within 10 m (33 ft) of any protected cave during fire ant surveys, routine maintenance, or any other management or monitoring activity or if biological investigations find any RIFA within any cave that has endangered invertebrates, all mounds within 10 m (33 ft) of that cave entrance should be treated within one week (USFWS 2011b). Staff conducting RIFA surveys as well as those conducting routine maintenance and other biological surveys on a Karst Feature Area should be trained to distinguish RIFA and their mounds from native ants and their mounds (USFWS 2011b).

Within 105 m of the entrance of any karst features that support listed invertebrates and/or SOCs, RIFA control must be restricted to the use of boiling water, which ensures protection from pesticides of the entire cave cricket foraging area (Taylor et al. 2005). In addition, RIFA bait treatments are not recommended outside of the cave cricket foraging area due to the fact that the baits can harm native ant species. For boiling water treatments, boiling or nearboiling water should be poured directly onto the mounds. Sufficient boiling water should be used that the mound collapses in on itself. This should typically be 1-4 gallons. These treatments should be conducted when the brood is high in the mound (typically on cool, cloudy days) to ensure that the queen(s) and larvae are likely to be near the top of the mound. During long periods of drought or cold, the queen(s) and larvae will most likely be deep within the mound, making them more difficult to eradicate (Vinson and Sorensen 1986). Mounds should not be disturbed before treatment as this will cause the ants to move the queen(s) and larvae to deeper locations within the mound or to a remote location (USFWS) 2011b). Small amounts (1-2 teaspoons) of detergent may be added to the boiling water, which helps the water penetrate the soil.

Passive management strategies should be implemented in conjunction with active management (boiling water treatments to mounds). Passive management strategies include: allowing woody vegetation to flourish and avoiding clearing of

native vegetation with the cave cricket foraging area to create a closed canopy which deters RIFA (RIFA's habitat preference is open/ disturbed habitat); controlling deer densities and feral hog populations, which can greatly increase woody growth and decrease soil disturbance; and not allowing public trails or picnic tables within the cave cricket foraging area.

Inspections will be made at cave sites during field visits for the presence of tawny crazy ant infestation. Managers will use current collection and reporting procedures of suspect infestations to confirm presence of new tawny crazy ant colonies, and if found, will work with the USFWS on control options.

Larger mammals, in particular raccoons, using cave features for shelter especially in and around urban areas can produce large amounts of scat inside the caves. The scat alters the nutrient content, especially nitrogen levels, within the cave ecosystem and can be detrimental to karst invertebrates (USFWS 2011e). Evidence of raccoon populations within caves should be monitored and populations controlled as needed.

6.2.3 Cave Gating and Fencing in BCP Caves and Bat Management

The need for a cave gate or protective fencing will be determined by each cave managing organization based on the following general criteria:

- In cases where caves are isolated (not near any neighborhoods), and/or with camouflaged entrances that do not appear to be a cave, no gates are warranted.
- 2. In situations where the cave has either a history of public access or is in near proximity to neighborhoods with a very obvious entrance, gating or fencing is recommended.
- 3. A gate or fence may also be necessary for liability reasons especially if the cave is vertical, unstable, or is a known "bad air" cave (USFWS 2011b).

Cave gates, where necessary, should be designed to permit normal airflow, water flow, and nutrient input, and should allow bat and small mammal (raccoon, opossum, fox, rodents, etc.) access (USFWS 2011b). Fences are an alternative to gating that may pose less interference with the nutrient regime and other

environmental factors (air and water movement). If the cave contains bats, then a fence may be more appropriate. The fence should be designed to be very difficult or close to impossible to climb over, and placed away from the cave entrance as far as possible (Sanders 1997 pers com). However, neither gates nor fences can prevent people from throwing toxic or other materials into a cave. Cave gates and fences may also serve to attract attention to an otherwise unknown cave which may encourage vandalism. Therefore, decisions about the need and desirability of gating or fencing BCP caves should be made on a case-by-case basis.

Bat gates should be installed on caves with suitable bat habitat. Prior to the construction of a cave gate, the cave should be evaluated for suitability as historic or current bat habitat. The criteria include historic bat use, numbers of bats currently using the cave, size of the entrance, size and arrangement of the interior rooms, surrounding habitat use, unavoidable disturbances from surrounding land use, and compatibility with other cave uses. Specialized gates will also be necessary for caves that receive large amounts of recharge. The design of bat gates should allow for access by the bats, by property managers, and by raccoons and small mammals, and should be as visually natural looking as possible. Information on bat gate design should be obtained from Bat International (www.bci.org) and/or Conservation the American Cave Conservation Association (www.cavern.org) to ensure there are no inadvertent impacts on karst invertebrates, bats or other species (USFWS 2011b).

Bat populations in caves should be monitored for potential effects of WNS on their numbers, and observations of multiple live or dead bats that exhibit signs of WNS should be reported immediately to the USFWS Austin Ecological Field Office. No bats are to be handled unless authorized to do so by the appropriate governmental agency (WNS Decontamination Team 2012). If WNS is discovered in the region in the future, BCP staff will follow appropriate decontamination procedures as outlined by the most recent National White-nose Syndrome Decontamination Protocol (WNS Decontamination Team 2012). Visitors from outside of central Texas or who have caved in Europe or any state where WNS has been suspected or confirmed, or researchers that request access to BCP caves must agree to adhere to the current WNS decontamination protocol prior to access or scientific research permit approval.

6.2.4 Physical Management

Cave areas should be protected from spills or contamination. The cave area is defined as the protection area designated by a hydrogeological investigation, or in the absence of a study, the area within 1/4-mile radius of the cave entrance. Coordination with USFWS is required if there are any possible contamination issues. Pesticides and fertilizers are prohibited from use within the area designated as needed for protection.

Electric power lines with transformers should be prohibited from critical cave areas because they could leak onto the ground or explode and adversely affect the cave fauna.

"Emergency Response Plans" (where needed) will be written in coordination with the Texas Commission on Environmental Quality (TCEQ), the COA Watershed Protection Department (WPD), and the Barton Springs/ Edwards Aquifer Conservation District (BS/EACD) for any cave near a pipeline or road where a major spill can occur. Most of the responsibilities for response will fall on these above agencies; however, creating such a plan before a spill may be critical to having the BCCP's interests represented and considered in a timely manner for protection of karst species.

No subsurface utility lines, roads or any other construction should enter or cross the cave area due to possible cave collapse, leakage from pipe corrosion, or related stresses. Altering and severing interstitial spaces negatively impacts and alters sensitive karst areas. Alteration of surface drainage patterns on BCP preserves without approval of USFWS will not be allowed.

BCP Partners will prevent dumping and vandalism at caves, and will remove trash from caves when encountered. When removing trash, BCP land managers will work to remove karst invertebrates from collected trash and return them to the cave.

6.2.5 Access Guidelines

In general, access to publicly-owned BCCP caves should be limited to necessary monitoring, management and research efforts that either directly benefit the endangered species or SOC or provide necessary maintenance (including RIFA control, gate maintenance, and insuring the security of the cave preserve). However, some publicly-owned BCCP caves have allowed public visitation since before the signing of the BCCP; this public access is considered to be grandfathered based on these prior allowances. Publicly owned BCCP caves with grandfathered status are: Airmen's Cave, District Park Cave, Goat Cave, Maple Run Cave, and Midnight Cave.

The Permit states that "all access to caves must be restricted to permits issued by the appropriate land management agency, based on an appropriate program in the land management plan for the preservation of the caves' ecosystem" (USFWS 1996a). BCP Partners will determine the type and amount of access at publicly-owned caves for the purposes of research, monitoring, or education, with priority focusing on adequate protection of karst species (See Tier II-A, Chapter XII). USFWS requires that anyone entering an endangered species cave without a 10(a)1(A) permit should be accompanied by someone who does have this permit.

USFWS (2011a) urges land managers to minimize access into caves due to impacts caused by visitation such as: increasing soil compaction, trash, and vandalism; scaring away trogloxenes; and direct mortality of cave organisms crushed by human disturbance. Human visitation may also disrupt cave ecosystems through introduction of non-native microorganisms, introduction of lint from clothing, increases in carbon dioxide, temperature, and nutrients, decreased humidity, and damage to speleothems (Hunter et al. 2004, Ilkner et al. 2007, Jablonsky et al. 1995, Lavoie and Northup 2005, Legatzki et al. 2011, Pulido-Bosch et al. 1997, U.S. Geological Survey 2013). Visitation impacts can be especially detrimental to low-energy caves (Gillieson 2011). Excessive or uncontrolled visitation may also endanger inexperienced people entering the

caves that are unguided, and/or lack proper safety devices and training, which could contribute to poor public opinion about caves, and can adversely affect the efforts of the BCCP. Because of these concerns, any BCCP caves that are open to the public through controlled, guided access should be accompanied by regular biological surveys of karst invertebrates as well as human visitation counts to assess impacts. Cave visitor impact monitoring may also be implemented to detect damage and guide visitor management. Trained volunteer cave monitors within Austin caving organizations could also play a vital part in the effort to protect BCP caves. These cavers can be a significant resource in cave management and will be allowed access to caves to assist with cave protection and education programs.

COA WPD staff is currently in the process of identifying new non-BCCP caves that will have the potential to reduce current levels of public access to BCCP caves while still providing valuable educational opportunities to the public.

If managing BCCP caves on private land, permission of the property owner or appropriate representative must always be obtained prior to entering. Good relationships with property owners of caves are valuable for promoting the goals of the BCCP, which includes securing the survival of rare and endangered karst species.

6.2.6 Public Education and Outreach

Education both for land management professionals and the general public should be implemented in order to raise awareness of cave conservation issues and encourage protection of caves and karst ecosystems. Education for BCP preserve managers, consultants, other professionals, and private landowners with BCP caves should be the immediate focus, which should include relaying up-to-date management strategies and monitoring efforts for determining and responding to the threats to karst ecosystems addressed above. Education for the general public should be a primary focus in the long term, to better inform citizens on the importance of protecting karst areas and how that protection also benefits them. For the purposes of this document, public education includes literature, curriculum, web media, interpretive kiosks, and guided surface and subsurface tours that can be made available for the general public, agencies, and individuals interested in learning more about karst areas and their inhabitants. Also included is educational media for cave managers and supporting staff, as well as the agencies involved with invertebrate species protection. A higher public awareness is an important step towards the recovery of the endangered cave invertebrates and continued preservation of karst species of concern.

6.3 MANAGEMENT COORDINATION

BCP partners will continue efforts to standardize management strategies and research/monitoring methods for all BCP caves based on best management practices.

BCP partners will attempt to work with and/or obtain landowner agreements with the following groups which are now protecting BCP caves (**Table 3**):

- TNC (one cave),
- TCMA (two caves),
- the Four Points cluster 10(a) agreement holder (five caves), and
- Canyon Creek as Section 7 agreement holder (one cave).

BCP partners will attempt to protect the remaining privately owned caves through acquisition, easements, or cave management agreements with the landowner. The precise location of some of these privately owned caves is currently unknown; therefore, the COA and TC should attempt to locate these caves in order to make a meaningful assessment.

Per BCCP Permit Conditions S2 and T2, if new karst features "are discovered with a significant diversity of troglobitic fauna, those karst features may be submitted to the Service for consideration for exchange with karst features identified for protection by the BCCP" (USFWS 1996b). In order to allow the Permit holders to implement these Permit conditions, COA and TC created a Cave Substitution Policy that provides a process that allows caves listed in the

BCCP permit to be substituted with other suitable caves in a manner that is transparent, science based, and consistent with the vision and intent established for BCCP. This policy includes a definition of "significant diversity of troglobitic fauna" as it applies to eligibility of a cave for substitution, and determines parameters that quantify preservation of environmental integrity for BCCP-listed caves and candidate substitution caves as it applies to management of caves. These defined criteria will be used in determining both the need to substitute a feature listed on the Permit as well as whether the substitution cave will adequately replace the previously identified BCCP cave or caves. The BCCP Cave Substitution Policy was adopted by the BCCP Coordinating Committee in August 2015, and is attached to this Land Management Plan as Appendix A.

The COA and TC will continue to monitor proposals for infrastructure projects that may impact BCP caves (see Management Handbook: Infrastructure).

BCP partners will continue to submit annual reports to the USFWS for all 62 caves detailing implementation of site specific management plans, cave acquisitions and agreements, research/monitoring results, and management actions and issues (USFWS 1996a).

7.0 MONITORING / RESEARCH

Monitoring Objectives include the following:

- Routine site inspections for signs of vandalism, unauthorized entry, trash dumping, presence of RIFA/tawny crazy ants, and damage to vegetation due to deer, feral hogs, or visitor off-trail use (USFWS 2011b).
- Verification of all BCP cave locations using established, systematic protocols. All BCP caves should also have interiors mapped using the most up-to-date survey methods available. When verifying cave locations, each site should be given a unique ID number using a tree tag and photos taken of each entrance.
- Baseline monitoring of cave species (listed and unlisted), cave crickets, vegetation, environmental conditions (in-cave and on the surface), RIFA, and mammals (USFWS 2011b).

 Monitoring of vegetation around karst features and within the features themselves for presence of feral hogs, deer, raccoons, etc. Monitoring will follow USFWS approved guidelines (USFWS 2011b).

Caves containing endangered and rare karst invertebrates on BCP properties will be monitored to determine long term trends in populations of cave organisms and overall cave conditions. All COA and TC owned BCCP caves will be surveyed Other BCP caves with ES/SOC will also be surveyed annually annually. dependent on staff availability. In addition, COA and TC identified 25 caves within Travis County managed by BCP partners that provide a more evenly distributed data set across cave clusters and karst faunal regions (KFRs). This new monitoring plan began in FY2011, with the number and frequency of karst faunal surveys and cricket counts synchronized among managing partners to better accommodate comparisons and determine trends. The goal of these changes to the cave monitoring program is to provide a clearer understanding of the species distribution and health of karst ecosystems across the BCP. Biomonitoring of the caves should follow methodology supported by USFWS to provide results that can be compared between caves throughout the region for better study and analysis.

All research, whether by BCP partners or outside researchers should not result in the "take" of an endangered species or in any way degrade endangered species habitat. All researchers must obtain approval from the land managers of BCP tracts being used for the research. If the proposed research involves endangered species the researchers must obtain a 10(a)1(A) permit from USFWS (USFWS 2011c). Land managers should also have potential researchers sign a standard form stating that they will abide by the rules of the BCP management plans or preserve rules.

The protocol for research and monitoring of cave fauna involves the use of 1-5 (depending on size of cave and logistics) pre-designated, permanent transects or zones per cave in which all living organisms encountered are identified and enumerated. Survey areas should be approximately 5 meters in length and span the width of the cave, or when possible, survey areas should occur in discreet

units of the cave such as a small room or an easily discernible section. Most importantly, the size and location of the survey area should remain constant during the course of the study. A non-toxic method of marking the transect boundaries (i.e. plastic flagging) may be necessary.

Ideally, each survey should be conducted by at least two people according to the cave's safety protocol. For each survey area, start and end time and the presence of trash or new vandalism will be recorded. Relative humidity and temperature will be recorded both outside the cave and at each transect or zone. Preferably, in order to standardize counts, the same observers will conduct all surveys. Typically observers start at opposite ends of the survey area and move toward each other while searching the cave floor, walls, ceilings, and beneath rocks for invertebrates. Any rocks that are lifted during the search will be replaced to their original position. Observers will be able to identify cave organisms to the nearest possible taxa (often genus or species), and will use a checklist of known invertebrates from the cave being surveyed. All data collected during cave surveys will be entered into the BCP Karst Database.

Any unknown invertebrates will be collected and identified by a karst invertebrate specialist. In caves containing endangered species, collecting should only occur in these caves with a special collecting permit obtained by USFWS. Observers should be extremely careful to not harm cave organisms while conducting surveys. All collected specimens should be deposited within the Texas Memorial Museum, or other reputable facility (USFWS 2011c). The date of deposition and collection number should also be recorded (USFWS 2011c). Additional procedures should continue to be developed to further define acceptable survey methods.

For caves that have controlled access, managers will keep records of every visit including information on: date, time, number of visitors, observations, temperature, humidity, etc.

Land managers will also monitor the entrances of caves containing endangered species at least twice yearly for anything that might harm the rare invertebrates

including presence of toxic substances, unauthorized use by recreational cavers, and surface disturbances which might have erosive potential or cause changes in surface drainage patterns. In addition, the interior of caves containing endangered species or SOC will be surveyed annually during dry, hot periods to check for RIFA infestation.

The overall health of caves can also be monitored by using semi-annual cave cricket exit counts. Cricket counts are done as they emerge from caves during good weather nights (i.e. not raining, warm etc.). The duration of the counts should remain constant (timed for two hours starting just after sunset). Additional information should be researched and incorporated into the methodology for conducting these cricket counts, as well as insight on how to relate survey data to cave health.

Groundwater and drip water samples should be collected to determine the impact of development on groundwater quality. Baseline sampling should be done in critical caves and springs. Tests should be done for geochemical mineral parameters as well as tests for heavy metals, organic chemicals and other likely pollutants. These tests should be done during development and for several years after development to determine if the groundwater and cave fauna are being adversely impacted by the changes in land use (Veni and Associates 1988). A list of parameters will be developed to standardize monitoring objectives. These should be listed in order of priority, should include sampling protocols, and should include a table of estimated and current year costs to assist landowners in budgeting management costs.

7.1 NEWLY DISCOVERED KARST FEATURES IN THE PERMIT AREA

If BCP Partners become aware of new cave and karst features (i.e. in projects submitted to these agencies during the development process), these features should be reported to the appropriate organizations such as USFWS, TCEQ, etc. Newly discovered karst features on BCP properties should be documented and species inventories done by BCP partners to provide information on potential new endangered species or SOC localities. When considering excavation of

newly discovered karst features for monitoring access, BCP land managers should consult cave excavation guidelines provided by USFWS (2011c).

7.2 RESEARCH NEEDS

There is currently insufficient information about some of the aspects of karst species and management of their habitat. In addition to the basic information listed above, BCP Partners should try to obtain information about the following topics and encourage research proposals and projects in these areas. This is not an exhaustive list, and research needs should be reviewed periodically. Research topics include:

- Cave Environments (humidity, temperature, airflow, and CO₂ concentrations). Increased airflow can cause the desiccation of cave passages. Also, the fluctuations of these abiotic conditions are not well documented in local caves and should be monitored to better understand potential impacts to karst invertebrates.
- Effects of opening or enlarging cave entrances. Excavating cave openings probably allows organic matter and nutrients to enter, and may enhance invertebrate diversity. For example, in Electromag Cave of Sun City, cave crickets became numerous after opening the entrance. However, it is possible that excavating these cave openings would enhance airflow and sunlight that may lead to drying of the cave. The general effect of opening caves probably results in returning the cave environment to pre-Colonial period conditions. This is because over grazing, agriculture, and other land-disturbance activities appear to have caused widespread filling of sinkhole depressions and cave entrances over the last few hundred years. The possible effect of opening or enlarging cave entrances requires further study. Criteria for determining the need for excavating karst features should be developed for the BCP, following cave excavation guidelines provided by USFWS (2011c).
- Delineating surface and sub-surface drainage basins to all BCP caves. Observations of flow inside caves and groundwater tracing should be used to better understand the water source for caves. Hauwert and Cowan (2013) provide methodology to adequately delineate the source area of cave drips and streams for achieving these goals.

- Life history studies. Information on the life history of karst invertebrate species on the BCCP permit is lacking and should he conducted. Life history studies that occur inside caves are best. Research of this type could potentially also be conducted at simulated cave environments, such as in the Austin Nature Center or in the Barton Springs Splash exhibit. Additionally there is a need to study habitat requirements of key trogloxene species such as cave crickets.
- Invasive species. RIFA, pill bugs, roaches, hothouse millipedes, and fleas can compete with or prey on other invertebrates. The degree of impact of these invasive species could be better understood. Attempts should be made to collect RIFA carrying prey to determine which species are most impacted. Understanding the effects of tawny crazy ants on karst ecosystems is also necessary. Finally, quantifying the effects of large amounts of scat in caves could be useful in understanding how this could attract non-cave adapted species such as roaches.
- Chemical impacts. Sampling and water-quality analysis of cave drips should be performed in urban areas, especially for pesticides, fertilizers, and metals. COA WPD tests groundwater for water-quality constituents from selected caves throughout the BCP. Local groundwater studies have found occurrences of lead, arsenic, petroleum hydrocarbons, and pesticides like bromacil and 4-nitrophenol in the groundwater under urban areas. Levels of hydrocarbon fumes have been documented in and near caves containing SOC (Get Down and Midnight caves), following a petroleum pipeline spill in 1986. The constituents of air in Travis County caves should be monitored periodically or in association with biological surveys.
- Aquatic life within the aquifer. Very little is known about life inside the aquifer in Travis County. Abundant diversity has been found in the Edwards Aquifer of the San Marcos to San Antonio area after investigation. Possible research could include: down-hole cameras and baited traps utilized in open bore wells; fine nets used to catch body parts in large capacity pumping wells; and surveys conducted in caves extending down to the water table. Efforts should be made to discover cave routes that extend to the water table, as these present tremendous opportunities to examine aquatic life.

- Cave cricket abundance as an indicator of cave ecosystem health. Cave cricket exit count data should be analyzed to determine trends. Studies on cave cricket foraging and surface habitat preferences should be conducted. Cave cricket survey methodology should be examined for improvements based on future scientific studies.
- Species identification. Efforts should be made to identify to species level yet undetermined troglobites in BCP caves, with special emphasis on species that may be identified as endangered or SOC as listed on the regional permit. Such examples include blind *Eidmannella* spiders, *Speodesmus* millipedes, *Rhadine* beetles, and Trichoniscidae isopods.
- Long-term trends in populations of cave organisms and overall cave conditions.
- Impacts on the species by recreational uses of caves (in caves with allowed access).
- Impacts of surface disturbances on karst species. Such disturbances may include reduced habitat area around the cave, erosion, changes in surface drainage patterns, and habitat restoration projects (mechanical clearing of vegetation and prescribed burns).
- Impacts from changes in surrounding land use. There is a need to better understand how development around cave areas may adversely impact groundwater, nutrient input, or the cave fauna themselves.

8.0 LITERATURE CITED

- Amos, B.B., and F.R. Gehlbach, 1988. Edwards Plateau Vegetation: Plant Ecological Studies in Central Texas. Baylor University Press, Waco, TX.
- Bayless, Todd, Travis County, Natural Resources and Environmental Quality, personal communication, 2013, 2015.
- Biological Advisory Team (BAT). 1990. Comprehensive report of the biological advisory Team. Austin, Texas.
- Campbell, J., Cokendolpher, J., Griswold, C., Ledford, J., Paquin, P. 2012.
 Systematics, conservation and morphology of the spider genus
 Tayshaneta (Araneae, Leptonetidae) in Central Texas Caves. Zookeys: (167): 1-102.
- Cokendolpher, James C., 2004. *Cicurina* spiders from caves in Bexar County, Texas (Araneae: Dictynidae). Texas Memorial Museum, Speleological Monographs, 6:13-58.
- Cokendolpher, James, Natural Science Research Laboratory, Museum of Texas Tech University, personal communication, 2008.
- Elliot, William R. 1992. Endangered and Rare Karst Species in Travis County, Texas: Options for the Balcones Canyonlands Conservation Plan.
- Elliott, William R. 1997. The Caves of the Balcones Canyonlands Conservation Plan, Travis County Texas, Travis County Natural Resources program.
- Elliott, William R. and J. R. Reddell. 1989. The status and range of five endangered arthropods from caves in the Austin, Texas, region. Appendix A in comprehensive report of the Biological Advisory Team, 1990.
- Gillieson, David S. 2011. Management of Caves. Chapter 6 in Karst Management, Phillip E. van Beynen (Ed.), p. 141-158.
- Hauwert, N., and Cowan, B. 2013, Delineating Source Areas To Cave Drips And Cave Streams In Austin Texas, USA: 13th Sinkhole Conference, Carlsbad, NM.

- Hedin, Marshall. 2015. High-stakes species delimitation in eyeless cave spiders (*Cicurina*, Dictynidae, Araneae) from central Texas. Molecular Ecology 24(2): 346-361.
- HNTB. 2005. Summary of Information for Assessing the Status of the Tooth Cave ground beetle (*Rhadine persephone,*), a report for USFWS by Casey Berkhouse.
- Hunter, A., D. Northup, C. Dahm, and P. Boston. 2004. Persistent coliform contamination in Lechuguilla Cave pools. Journal of Cave and Karst Studies 66(3):102-110.
- Ikner, L., R. Toomey, G. Nolan, J. Neilson, B. Pryor, and R. Maier. 2007. Cultural microbial diversity and the impact of tourism in Kartchner Caverns, Arizona. Microbial Ecol. 53:30-42.
- Intergovernmental Panel on Climate Change (IPCC). 2007. Climate change 2007: synthesis report, summary for policymakers. Intergovernmental Panel on Climate Change, Fourth Assessment Report. Released on 17 November 2007. 23 pp. Available from: http://www.ipcc.ch/pdf/assessment report/ar4/syr/ar4_syr_spm.pdf.
- Jablonsky, P., S. Kraemer, and B. Yett. 1995. Lint in caves. Proceedings of the 1993 Cave Management Symposium. Carlsbad Caverns, New Mexico. p. 73-86.
- Lavoie, K. and D. Northup. 2005. Bacteria as indicators of human impact in caves. Proceedings of the 2005 National Cave and Karst Management Symposium. Albany, New York. p. 40-47.
- Legatzki, Antje, et al. "Bacterial and archaeal community structure of two adjacent calcite speleothems in Kartchner Caverns, Arizona, USA." Geomicrobiology Journal 28.2 (2011): 99-117.
- Ledford, Joel, Department of Entomology, California Academy of Sciences, personal communication, 2010.

- Meyers, Jason M. 2008. Identification, Distribution, and Control of an Invasive Pest Ant, *Paratrechina* sp. (Hymenoptera:Formicidae), in Texas. PhD. Dissertation. Texas A and M University.
- Muchmore, William B. 2001. Review of the genus *Tartarocreagris*, with descriptions of new species (Pseudoscorpionida: Neobisiidae). Texas Memorial Museum, Speleological Monographs, 5:57-72.
- Paquin, P. and M. Hedin. 2005. Genetic and morphological analysis of species limits in *Cicurina* spiders (Araneae, Dictynidae) from southern Travis and northern Hays counties (TX), with emphasis on *Cicurina cueva* Gertsch and relatives. Special report for the Department of Interior United States Fish & Wildlife Service Contract No. 201814G959. Revised version10 May 2005.
- Paquin P., Dupérré, N., Cokendolpher J., White, K.and M. Hedin. 2008. The fundamental importance of taxonomy in conservationbiology: the case of the eyeless *Cicurina bandida* (Araneae:Dictynidae) of central Texas, including new synonyms and the description of the male of the species. Invertebrate Systematics 22: 139–149.
- Pulido-Bosch, A., W. Martin-Rosales, M. Lopez-Chicano, C. Rodriguez-Navarro, and A. Vallejos. 1997. Human impacts in a touristic karstic cave (Aracena, Spain). Environmental Geology 31:142-149.
- Reddell, James, Texas System of Natural Laboratories (TSNL), Texas Cave Management Association (TCMA), personal communication, 1998, 2004, 2005.
- Reddell, James, Texas Memorial Museum, University of Texas, personal communication, 2010.
- Sanders, Mark, City of Austin, Austin Water Utility, BCP Program, personal communication, 1997, 2013, 2015.
- Taylor, S.J., J.K. Krejca, and M.L. Denight. 2005. Foraging range and habitat use of Ceuthophilus secretus (Orthoptera: Rhaphidophoridae), a key

trogloxene in central Texas cave communities. American Midland Naturalist 154: 97-114.

- Texas Memorial Museum (TMM). 2007. TEXBIO database, The University of Texas at Austin.
- U.S. Fish and Wildlife Service. 1994. Endangered Karst Invertebrates (Travis and Williamson Counties): Recovery Plan.
- U. S. Fish and Wildlife Service, 1996a. Final Environmental Impact Statement/Habitat Conservation Plan for Proposed Issuance of a Permit to Allow Incidental Take of the Golden-cheeked Warbler, Black-capped Vireo, and Six Karst Invertebrates in Travis County, Texas. U. S. Fish and Wildlife Service, Albuquerque, N. M.
- U. S. Fish and Wildlife Service. 1996b. Federal Fish and Wildlife Permit No. TE-788841-2.
- U.S. Fish and Wildlife Service. 2008. Tooth Cave Ground Beetle (*Rhadine Persephone*) 5-Year Review: Summary and Evaluation. Austin Ecological Services Field Office, Austin, TX.
- U.S. Fish and Wildlife Service. 2009a. Bone Cave Harvestman (*Texella reyesi*) 5-Year Review: Summary and Evaluation. Austin Ecological Services Field Office, Austin, TX.
- U.S. Fish and Wildlife Service. 2009b. Tooth Cave Spider (*Neoleptoneta myopica*), Kretschmarr Cave Mold Beetle (*Texamaurops reddelli*), and Tooth Cave Pseudoscorpion (*Tartarocreagris texana*) 5-Year Review: Summary and Evaluation. Austin Ecological Services Field Office, Austin, TX.
- U.S. Fish and Wildlife Service. 2009c. Bee Creek Cave Harvestman (*Texella reddelli*) 5-Year Review: Summary and Evaluation. Austin Ecological Services Field Office, Austin, TX.
- U.S. Fish and Wildlife Service. 2011a. Bexar County Karst Invertebrates Recovery Plan. USFWS Southwest Region, Albuquerque, NM.

- U.S. Fish and Wildlife Service. 2011b. Karst Preserve Management and Monitoring Recommendations. Austin Ecological Services Field Office, Austin, TX.
- U.S. Fish and Wildlife Service. 2011c. Section 10(a)(1)(A) Karst Invertebrate Survey Requirements for Conducting Presence/Absence Surveys for Endangered Karst Invertebrates in Central Texas. Fish and Wildlife Service, 10711 Burnet Road, Suite 200, Austin, Texas 78758. September 8, 2011. 21 pp.
- U.S. Fish and Wildlife Service. 2011d. A National Plan for Assisting States, Federal Agencies, and Tribes in Managing White-Nose Syndrome in Bats.US Fish & Wildlife Service, 300 Westgate Center Drive, Hadley, MA.
- U.S. Fish and Wildlife Service. 2011e. Karst Invertebrate Habitat Requirements. Austin Ecological Services Field Office, Austin, TX.
- U.S. Fish and Wildlife Service. 2012. Karst Preserve Design Recommendations. Austin Ecological Services Field Office, Austin, TX.
- U.S. Fish and Wildlife Service. 2014a. Service Determines that the Warton's Cave Meshweaver does not Warrant Protection Under the Endangered Species Act. U.S Fish and Wildlife Service Southwest Region News Release, August 13, 2014.
- U.S. Fish and Wildlife Service. 2014b. White Nose Syndrome Map Suspected and Confirmed Occurrences. WNS Coordination Node (online resource). <u>http://www.whitenosesyndrome.org/resource/white-nose-syndrome-</u> <u>infographic-updated-april-11-2014</u> (accessed 22 April 2014).
- U.S. Geological Survey. 2013. Threats to cave ecosystems. Retrieved July 2013 from http://sbsc.wr.usqs.gov/cprs/research/projects/caves/threats.asp.
- Veni and Associates. 1988. Hydrologic investigation of the Jollyville Plateau karst, Travis County, Texas. U.S. Fish and Wildlife Service.

Vinson, S.B. and A.A. Sorensen. 1986. Imported fire ants: Life history and impact. Texas Dept. of Agriculture, Austin, TX. 28p.

Watson, Cyndee, U.S. Fish and Wildlife Service, Austin Ecological Services Field Office, personal communication, 2015.

WNS Decontamination Team. 2012. National White-Nose Syndrome Decontamination Protocol - Version 06.25.2012. http://whitenosesyndrome.org/sites/default/files/resource/national_wns_re vise_final_6.25.12.pdf

- Zara Environmental, LLC. 2007a. Report of Excavation and Biology of Down Dip Sink, Water Treatment Plant 4 Bull Creek Site, Travis County, Texas. Report for Carollo Engineers, Austin TX.
- Zara Environmental, LLC. 2007b. Report of Excavation and Biology of Garden Hoe Cave and Canopy Joint Sink, Water Treatment Plant 4 Bull Creek Site, Travis County, Texas. Report for Carollo Engineers, Austin TX.
- Zara Environmental, LLC. 2008. Excavation and Biology of Caves and Karst Features on the Lucas Tract, Travis County, Texas. Report for Carollo Engineers, Austin TX.
- Zara Environmental, LLC. 2010. Population Status of Karst Invertebrates in the Balcones Canyonlands Preserve. Report for Weston Solutions, Inc., Austin TX.

APPENDIX A

BCCP CAVE SUBSTITUTION POLICY



Balcones Canyonlands Conservation Plan Coordinating Committee Cave Substitution Policy

Adopted: 8/2//15

Precinct 3 County Commissioner Gerald Daugherty

District 7 City Council Member Leslie Pool

Acknowledged:

William A. Conrad, Coordinating Committee Secretary

Purpose

Provide a process that that will allow the Balcones Canyonlands Conservation Plan (BCCP) Coordinating Committee to implement conditions in the BCCP Endangered Species Permit (ES 788841-2 and future revisions, amended permits, or subsequent permits) that allow caves listed in the permit to be substituted with other suitable caves in a manner that is transparent, science based, and consistent with the vision and intent established for BCCP

BCCP Permit Conditions

BCCP Permit (TE 788841-2) Condition S2 [related to Endangered Species covered by the permit]: If during investigations for development of a tract, karst features are discovered with a significant diversity of troglobitic fauna, those karst features may be submitted to the Service for consideration for exchange with karst features identified for protection by the BCCP. The determination of "significant diversity" will be made by the permit applicants and the Service, in association with karst experts. The inclusion of such a karst feature would not increase the number of caves to be protected by the BCCP, but would result in the new feature replacing a previously identified cave or caves.

BCCP Permit (TE 788841-2) Condition T2 [related to Species of Concern covered by the permit]: If during investigations for development of a tract, karst features are discovered with a significant diversity of troglobitic fauna, those karst features may be submitted to the Service for consideration for exchange with karst features identified for protection by the BCCP. The determination of "significant diversity" will be made by the permit applicants and the Service, in association with karst experts. The inclusion of such a karst feature would not increase the number of caves to be protected by the BCCP, but would result in the new feature replacing a previously identified cave or caves.

Roles and Responsibilities

BCCP Coordinating Committee - was created to implement the BCCP - Shared Vison. The BCCP Interlocal Cooperation Agreement requires that the Coordinating Committee carry out the BCCP Shared Vision as Authorized by the BCCP federal permit. It may adopt policies recommended by its Secretary. They are responsible for adopting this policy, providing public involvement with respect to its implementation, and for making decisions and taking action as provided by this policy including initial approval that would trigger actions to initiate a minor permit amendment.

Permit Covered Governing Bodies - as provided in the BCCP Interlocal Cooperation Agreement include Austin City Council and Travis County Commissioners Court. They are responsible for providing additional opportunities for public involvement and reviewing the Coordinating Committee's decisions to substitute caves for those covered in the permit and for taking action as they deem appropriate as provided for in the BCCP Interlocal Cooperation Agreement for permit amendments.

U.S. Fish and Wildlife Service (Service) - is the federal agency authorized by the Endangered Species Act to issue, suspend, and revoke incidental take permits in accordance with Section 10(a)(1)(B) of the Endangered Species Act and its implementing regulations, policy and guidance. They issued permit TE-788841-2 based on the March 1996 Habitat Conservation Plan and Final Environmental Impact Statement. Under their authorities, they are responsible for reviewing and approving any requests to amend this permit. The Service's role is to advise the BCCP on matters related to permit compliance and Fish and Wildlife Service processes and procedures at the earliest possible opportunity.

Third Parties - many caves identified for protection in BCCP are on property owned by third parties not bound to BCCP. They are not required to coordinate with or seek approval from BCCP before taking action that may affect a BCCP listed cave. Third parties may offer cave protection to BCCP for permit covered caves or for caves that might be considered for substitution by BCCP.

Cave Substitution Process

Background

The regional Endangered Species Act Section 10(a)(I)(B) permit (TE 788841-2), also known as the Balcones Canyonlands Conservation Plan (BCCP), requires the Permit holders (City of Austin and Travis County) to acquire, protect, and ensure management that preserves the environmental integrity of 62 listed caves protecting populations of six endangered karst invertebrates and 25 karst species of concern (SOC). This Permit "is subject to compliance with, and implementation of, the terms and conditions of the Environmental Impact Statement and Habitat Conservation Plan" (EIS/HCP) as well as all specific conditions contained in the Permit itself (USFWS 1996a).

One such condition described in the Permit states that "if during investigations for development of a tract, karst features are discovered with a significant diversity of troglobitic fauna, those karst features may be submitted to the Service for consideration for exchange with caves identified for protection by the BCCP. The determination of 'significant diversity' will be made by the permit applicants and the Service, in association with karst experts. The inclusion of such a karst feature would not increase the number of caves to be protected by the BCCP, but would result in the new feature replacing a previously identified cave or caves" (USFWS 1996a). In order to meet the terms and conditions of the U.S. Fish and Wildlife Service (USFWS) Permit, Permit holders determined a need to define "significant diversity of troglobitic fauna" as it applies to eligibility of a cave for substitution, and determine parameters that quantify preservation of "environmental integrity" for BCCP-listed caves and candidate substitution caves as it applies to management of caves. These defined criteria will be used in determining both the need to substitute a feature listed on the Permit as well as whether the substitution cave will adequately replace the previously identified BCCP cave or caves. These criteria are not intended to evaluate whether a BCCP-listed caves has met Permit compliance, but rather only to evaluate caves for substitution. Following Permit conditions, a group of karst experts, USFWS staff, and Permit holder staff collaborated on these criteria as members of the BCCP Scientific Advisory Committee Karst Sub-committee (chair: Dr. Nico Hauwert).

Significant Troglobitic Diversity as Applied to Conservation of Karst Species

Due to the predicted loss of the vast majority of potential karst habitat allowed by the BCCP, the EIS/HCP states that "the adequacy of the plan is contingent upon full implementation of the acquisition and management strategies detailed in the BCCP", which includes caves named as specific localities for the six endangered karst invertebrates and 25 BCCP-listed karst SOCs (USFWS 1996b). The EIS/HCP also stresses that given the fact that several BCCP karst SOCs were known from only a few caves when the plan was written, the loss of even one BCCP-listed cave could result in a major reduction to the species' population (USFWS 1996b).

However, the EIS/HCP acknowledges that although the BCCP was designed to protect most known localities of endangered karst invertebrates and karst SOCs at the time of permit issuance, "the possibility remains that features may be found that provide habitat for listed species or other equally rare karst invertebrates", and there is a "high probability that other new rare species will be described from Travis County in the future" (USFWS 1996b). The Permit provides for these types of new discoveries to be considered substitutions for BCCP-listed caves if such karst features provide a "significant diversity of troglobitic fauna" (USFWS 1996a).

In order to adhere to the protection strategy in the EIS/HCP and Permit for listed karst invertebrates, as well as ensuring Permit holders receive the "No Surprises" guarantee for protecting the 25 karst SOCs, a karst feature considered for exchange with a BCCP-listed cave must consider those species for which the BCCP cave was designated to protect. However, the EIS/HCP also guides Permit holders to attempt to protect newly discovered karst features that provide habitat for other equally rare karst invertebrates (USFWS 1996b).

The USFWS Biological Opinion also states that the BCCP "identifies an option that establishes a process that allows any newly discovered cave to be protected in the place of a less biologically significant cave currently identified for protection" (USFWS 1996c).

Incorporating this guidance from the EIS/HCP and Biological Opinion, the determination of a replacement cave's significant biological diversity will consider several factors that include species composition, diversity, and abundance, as well as the cave's location and ecological benefits. See Methodology for Assessing Significant Diversity and Environmental Integrity of BCCP Caves and Potential Substitution Caves for specifics on these factors and methodology used for determining significant troglobitic diversity of karst features.

Environmental Integrity as Applied to Karst and Caves

The EIS/HCP states that for a karst fauna area to be considered protected, it must "contain a large enough expanse of continuous karst and surface area to maintain the integrity of the karst ecosystem on which each species depends." The EIS/HCP also provides protection criteria to meet this goal, stating that "the size and configuration of each karst fauna area must be adequate to maintain moist, humid conditions, air flow, and stable temperatures in the air-filled voids; maintain an adequate nutrient supply; prevent contamination of surface and groundwater entering the ecosystem; prevent or control the invasion of exotic species, such as fire ants; and allow for movement of the karst fauna and nutrients through the interstitium between karst features" (USFWS 1996b).

The EIS/HCP states that, "in most instances, this will entail protecting the entire surface and sub-surface drainage area of each cave and enough of the surface vegetation community to support small animals and buffer against fire ant infestations" (USFWS 1996b).

Although the 1996 EIS/HCP does not provide a quantifiable area for protection of the surface vegetation community, it does address the need for this information by stating that the delineation of appropriate boundaries for individual cave preserves will require additional studies to determine the surface area necessary to maintain the biological resources important to the cave (USFWS 1996b).

Research and information needs such as this were also outlined in the Recovery Plan for Endangered Karst Invertebrates in Travis and Williamson Counties, Texas (1994) as one of four major recovery actions; the EIS/HCP reiterates that the effective enactment of this and other recovery actions are necessary "to assure that the implementation of the BCCP has no negative impact on the population viability of the endangered karst invertebrates" (USFWS 1996b).

This recovery action was met with the completion of USFWS Karst Preserve Design Recommendations in 2012, which quantifies the protection criteria quoted above from the BCCP EIS/HCP, and provides specific preserve components for configuring karst preserves that maintain environmental integrity of the karst invertebrate locations and ecosystems they are designed to protect.

According to USFWS' Karst Preserve Design Recommendations, in addition to protecting the entire surface and sub-surface drainage areas, preserve components which maintain the cave's environmental integrity should include: the cave cricket foraging area; a preserve configuration of at least 40 acres that protects the surface plant and animal communities and ensures that the cave footprint is over 105 meters from the nearest hard edge; and is free of incompatible forms of land use and sources of contamination (USFWS 2012).

These recommendations (USFWS 2012) also reiterate the need for karst preserves to be protected and management assured through acquisition or formal management agreements, which is also a requirement of the Permit and EIS/HCP (USFWS 1996a, USFWS 1996b).

Additional preserve components meeting these objectives and methods for quantifying and evaluating these factors are described in Methodology for Assessing Significant Diversity and Environmental Integrity of BCCP Caves and Potential Substitution Caves.

Caves submitted as substitution caves for BCCP-listed caves will be assessed for their environmental integrity using these factors, and measured against the environmental integrity assessment of the cave or caves suggested for replacement.

Only replacement caves with sufficient environmental integrity and significant diversity of troglobitic diversity, and equivalent to or superior to the BCCP cave it has been submitted to replace will be used as an adequate substitution. See Methodology for Assessing Significant Diversity and Environmental Integrity of BCCP Caves and Potential Substitution Caves for methodology on factors for determining environmental integrity of karst features.

Literature Cited

U. S. Fish and Wildlife Service. 1994. Recovery Plan for Endangered Karst Invertebrates in Travis and Williamson Counties, Texas. Albuquerque, New Mexico.

U. S. Fish and Wildlife Service. 1996a. Federal Fish and Wildlife Permit No. TE-788841-2.

U. S. Fish and Wildlife Service. 1996b. Final Environmental Impact Statement /Habitat Conservation Plan for Proposed Issuance of a Permit to Allow Incidental Take of the Goldencheeked Warbler, Black-capped Vireo, and Six Karst Invertebrates in Travis County, Texas.

U. S. Fish and Wildlife Service. 1996c. Biological Opinion for the Issuance of a Section 10(a) (I)(B) Permit for the Balcones Canyonlands Conservation Plan. Albuquerque, New Mexico.

U. S. Fish and Wildlife Service. 2012. Karst Preserve Design Recommendations. Austin Ecological Services Field Office. Austin, Texas.

Methodology for Assessing Significant Diversity and Environmental Integrity of BCCP Caves and Potential Substitution Caves

The following methodology describes criteria that will: 1) define significant diversity of troglobitic fauna in caves considered for replacement of BCCP-listed caves and 2) determine protection measures that quantify preservation of environmental integrity for BCCP-listed caves and candidate substitution caves.

These criteria will be used in determining both the need to substitute a feature listed on the Permit, such as when a cave ecosystem has been significantly damaged or destroyed or if Permit holders have been unable to secure adequate protections for a cave, as well as whether the substitution cave will adequately replace the previously identified cave or caves.

Caves submitted as substitution for BCCP-listed caves will be assessed for equal or superior significant diversity of troglobitic fauna and environmental integrity using the factors below, and measured against the assessment of the cave or caves suggested for replacement.

Evaluation to Determine Substitution Need and Suitability of Replacement Caves

Caves submitted as substitution caves for BCCP-listed caves will be assessed for whether they meet objectives for significant diversity of troglobitic fauna and environmental integrity using the

factors detailed below, and measured against equal assessments of the cave or caves suggested for replacement.

Only replacement caves meeting significant diversity requirements and with sufficient environmental integrity equal or superior to the BCCP cave it has been submitted to replace will be used as an adequate substitution.

Essential to this evaluation is that the replacement cave(s) be a confirmed locality for the same federally listed karst invertebrate(s) and/or karst species of concern (SOC) as the BCCP-listed cave designated for substitution.

Evaluations for BCCP-listed caves and candidate replacement cave(s) will be made using the most up-to-date research and karst preserve design recommendations available at the time of the assessment. If additional research valuable to this evaluation process becomes available in the future, the BCCP Scientific Advisory Committee's karst sub-committee will review the new information and incorporate or revise assessment factors below if deemed appropriate.

This document is not intended to be a precise rating system or contain a complete scoring rubric, but rather serves as a comprehensive list of the data that would be ideal to have in hand to evaluate cave substitution.

Caves are not easily comparable in terms of biology, ecosystem health, and value to preserve strategies. Each situation is different, and it is impossible to anticipate the variety of issues that may arise when comparing two caves.

The purpose of this document is to provide the evaluation team with a list of all reasonably measured factors relevant to the decision for approving a cave for substitution.

This document is strictly designed for the cave substitution process and not intended to be used to evaluate whether a BCCP-listed cave's protections are compliant with the Permit.

Evaluations for proposed substitutions will occur on a case by case basis, which includes determining if sufficient data are available to evaluate both BCCP-listed caves proposed for substitution and their candidate replacement cave(s).

If there are too many unknowns or assumptions about either cave, evaluators are allowed to reject the substitution proposal until the proposer fills in more of the dataset, up to the discretion of the evaluation team.

Proposers allowing cave access to evaluators may provide one option for obtaining missing evaluation data.

Factors for Determining Significant Diversity of Troglobitic Fauna for a Candidate Substitution Cave

Candidate replacement caves will be compared with BCCP caves designated for substitution and will only be accepted as replacements if the following conditions are met:

- 1. Replacement cave has similar or greater species composition in relation to target species (federally listed taxa or karst SOCs), as determined by the following factors:
 - a. The replacement cave must be a confirmed locality for the same federally listed karst invertebrate(s) as the BCCP cave it will be replace.

- b. For BCCP caves containing one of the 25 karst SOC, the replacement cave must contain the same SOC(s) as the BCCP cave it will replace.
 - i. Exception: If the BCCP cave does not contain any of the 25 karst SOC (Talus Spring Cave), then the replacement cave must contain either: 1) one or more karst SOC listed on the Permit or 2) one or more troglobitic species of similar taxa to the SOCs listed on the Permit considered to be at least as rare as the BCCP-listed SOCs.

Rare karst invertebrates not listed as SOCs on the Permit will be evaluated using information from the BCP Karst Database, Texas Memorial Museum's TexBio Database, Texas Park and Wildlife Department's Species of Greatest Conservation Need (SGCN) list rankings (TPWD 2011), and NatureServe's Conservation Status Assessments (Master et al. 2012) to quantify significance of the species in terms of rarity and need for protection.

- 2. Replacement cave has similar or greater overall troglobitic karst invertebrate species diversity, as determined by the following factors:
 - a. Demonstrate through repeated biological surveys that the replacement cave has greater or equal diversity; for example, by graphing the number of troglobitic taxa seen on each visit and noting those previously seen vs. new occurrences. Karst invertebrate surveys should follow survey methodology described in USFWS 2014b, which explains that in order to assess presence/absence of endangered karst invertebrates with a high level of confidence, caves should be surveyed at least 14 times.
 - b. Evaluate whether the caves in consideration have been thoroughly measured in terms of diversity. Since many karst species are rare and poorly studied, problems with detection and taxonomy hamper creating a complete list. Evaluate and explain the status of the diversity list for the cave(s).
 - c. Evaluate the numbers of troglobitic taxa vs. other taxa (troglophiles, trogloxenes, or accidentals). In some cases the cave entrance has great diversity, but the deep cave community structure is limited.
 - d. Additional rare karst invertebrates not listed as SOCs on the Permit will be evaluated using information from the BCP Karst Database, Texas Memorial Museum's TexBio Database, Texas Park and Wildlife Department's SGCN list rankings (TPWD 2011), and NatureServe's Conservation Status Assessments (Master et al. 2012) to quantify significance of the species in terms of rarity and need for protection.
 - e. Consider non-troglobitic karst species, which rank high on TPWD's SGCN list and NatureServe's Conservation Status Assessments list that could be affected/protected by the substitution (ex: bats, salamanders), as contribution to the overall biological diversity of the cave being considered for substitution.
- 3. Replacement cave has similar or greater Permit-listed species abundance, as determined by the following factors:
 - a. Demonstrate through repeated biological surveys the relative abundance of taxa on the cave's species list. With well-delineated in-cave survey methodology, it should be clear where the rare species are found within the cave, and how many are typically seen in a visit. If collection methods are not performed in a uniform fashion, results may not be comparable among sites or within a site on different survey days; this should be explained or accounted for in the evaluation. Karst invertebrate surveys should follow survey methodology described in USFWS 2014b.

- b. If the entire cave isn't inventoried during each visit, then an estimate of the nonsurveyed area would help determine the total potential of the cave to support a healthy population. This estimate will account for the fact that, on average, larger caves have more habitat available and a greater diversity of habitat, thus having a greater diversity and abundance of fauna (Schneider and Culver 2004).
- 4. Replacement cave's location is within the same karst fauna region as defined by Veni (1992) or future USFWS-approved revisions of the KFRs.
- 5. Replacement cave's location is within the same BCCP-protected cave cluster (Northwood, McNeil, or Four Points). Note: only applicable if BCCP cave to be substituted is within one of these cave clusters. This requirement ensures that the replacement cave is contributing to a Karst Fauna Area that helps meet the recovery criteria for the federally listed karst invertebrates in the BCCP cave to be substituted.

Factors for Determining Environmental Integrity of BCCP Listed Caves and Potential Replacement Caves

The following protection criteria, largely based on USFWS Karst Preserve Design Recommendations (2012), will be used to quantify the environmental integrity of BCCP caves and candidate substitution caves for determining both the need for substitution of a BCCP cave and adequate replacement by a candidate substitution cave.

Ideally preferred protection goals are also described for each factor to guide evaluation assessments.

A. Karst feature surface area protection measures:

- 1. Percent of cave footprint within protected area:
 - a. Determined by GIS spatial analysis and use of footprint digitized from cave map to quantify percent protected.
 - b. Protected area lands owned or acquired by the Permit holders (City of Austin and Travis County) or BCP managing partners that are managed for protection of the cave or caves, or lands that have formal management agreements with the Permit holders as described in S-4 and T-4 of the Permit (USFWS 1996a).
 - c. Ideally preferred protection goal: 100 percent of cave footprint is within protected area (USFWS 2012).
- 2. Distance of cave footprint to nearest preserve edge:
 - a. Determined by GIS spatial analysis by calculating the distance of edge of the digitized cave footprint to nearest preserve boundary.
 - b. Edge: defined as the cave preserve's property boundary and/or where impervious cover dissects the natural area surrounding the karst feature, such as paved roads or urban development areas detrimental to surface protection efforts.
 - c. Ideally preferred protection goal: footprint is as near to the center of the protected area as possible, and at least 105 m from the preserve edge (USFWS 2012).
- 3. Percent of surface drainage within protected area:
- a. Determined by GIS spatial analysis by quantifying percent of delineated surface drainage basin that is within protected area(s).

- b. Surface drainage basins will be conservatively over-estimated with high confidence by licensed geologists performing hydrogeologic studies of caves using methods described in Veni 2003, Hauwert et al. 2005, Hauwert 2009, or other methods approved by the BCCP Scientific Advisory Committee's karst subcommittee.
- c. Where surface drainage basin delineations are unable to be performed due to denied access, this analysis will be performed based on the area draining to the cave entrance using available topographic and cave map data.
- d. Ideally preferred protection goal: the entire surface drainage basin is within the protected area (USFWS 2012).
- 4. Percent of subsurface drainage basin within protected area:
 - a. Determined by GIS spatial analysis by quantifying percent of delineated subsurface drainage basin that is within protected area(s).
 - b. Subsurface drainage basins will be conservatively over-estimated with high confidence by licensed geologists performing hydrogeologic studies of caves using methods described in Hauwert and Cowan 2013, Veni 2003, or other methods accepted by the SAC karst sub-committee.
 - c. Where subsurface drainage basin delineations are unable to be performed due to denied access, this analysis will be performed using a delineation made by the contour level at the bottom of the cave, as required by S-3 and T-3 of the Permit (USFWS 1996a).
 - d. Ideally preferred protection goal: the entire subsurface drainage basin is within the protected area (USFWS 2012).
- 5. Percent of cave cricket foraging area (105 meters from cave footprint) within protected area:
 - a. Determined by GIS spatial analysis by creating a 105m buffer area around the cave's footprint digitized from its cave map, and quantifying percent of this buffer area that is within protected area(s).
 - b. Ideally preferred protection goal: 100 percent of cave cricket foraging area is within the protected area (USFWS 2012).
 - c. As an alternative to assuming a 105m buffer, site-specific cave cricket surveys could be performed in order to determine the foraging area around a specific cave. Methods should include an adequate survey area and effort during appropriate season and over enough nights to capture the large diversity of exit and foraging patterns known for *Ceuthophilus* spp. Taylor et al. (2005) and Zara Environmental (2013, 2014) give examples of methods used for site specific cave cricket foraging studies in Texas.
- 6. Preserve tract size:
 - a. Determined by GIS spatial analysis. In cases where BCP or other preserve tracts are adjoining each other, all connected interior preserve tract boundaries will be dissolved to account for connectivity to all preserve areas. Cave preserve tract delineations also cease at hard edges such as paved roads or impervious cover detrimental to surface protection efforts.
 - b. NOTE: This environmental factor is extremely important when determining environmental integrity of a candidate replacement cave. A large, intact tract has ecological stability and natural buffers that are difficult if not impossible to create artificially or manage successfully. Large preserves protect the quality of native

surface plant, arthropod, and animal communities necessary for adequate nutrient input (USFWS 2012). Large preserves are also more resilient and typically support caves that need less active management (USFWS 2012). Large preserves with contiguous karst areas have abundant mesocavernous zones which are likely to support immeasurable populations of rare troglobites. Having naturally resilient preserves also makes them less sensitive to problems associated with loss of funding and staffing resources that may happen over time.

- c. USFWS has chosen preserve size as a critical indicator in determining quality of a karst preserve (USFWS 2012). Ideally preferred protection goals include a preserve size of at least 100 acres to be considered a high quality preserve, or at least 40 acres to be considered a medium quality preserve (USFWS 2012).
- 7. Net gain in protected land for BCP:
 - a. Determined by subtracting acreage of BCCP cave's protected area from the replacement cave's preserve tract size.
 - b. Ideally preferred protection goal: Cave preserves with larger protected areas will be favored due to benefits described above in item 6, preserve tract size.
- 8. Shape of protected area:
 - a. Subjective determination using map that shows cave's location within delineated preserve area boundaries.
 - b. Ideally preferred protection goal: USFWS 2012 defines preserves that are circular in shape and/or are connected to other preserves as an ideally preferred protection measure, along with the cave or caves being as near to the center of the preserve area as possible to reduce edge effects.
- 9. Landscape mosaic of protected area (% woodland/grassland):
 - a. Determined by GIS spatial analysis using NAIP aerial imagery and/or LIDAR data to classify landscape components in the protected areas within 100 acres of cave.
 - b. Proper landscape mosaic helps to ensure the quality of the native surface plant, arthropod, and animal communities, beneficial to the cave's nutrient input (USFWS 2012).
 - c. Ideally preferred protection goal: according to USFWS (2012), cave preserve areas should include ≥80% woodland to 10% grassland mosaic.
- 10. Number of adjacent karst features within protected area:
 - a. Quantified by performing karst feature surveys in protected area within a 100 acre radius of cave which includes the surface and subsurface drainage basins, following recommendations in USFWS 2014b.
 - b. Ideally preferred protection goal: cave preserves should be designed to protect as many caves or karst features as possible to support nutrient input from cave crickets (USFWS 2012).

11. Incompatible land use/fragmentation:

- a. Subjective determination using aerial map that demonstrates the cave's location and incompatible forms of land use within delineated preserve area boundaries.
- b. Incompatible forms of land use within the delineated karst preserve itself such as paved roads, impervious cover, livestock, water retention ponds, or hiking and

biking trails should also be documented for consideration of this factor (USFWS 2012). Also describe adjacent land use outside of the preserve including developments, roads, impervious cover, etc.

- c. Ideally preferred protection goals: There should be no paved roads, development, impervious cover or other structures that result in permanent habitat loss within the cave's protected area (USFWS 2012). Protected areas should also not include trails or picnic tables inside the cave cricket foraging area, the surface or subsurface drainage basin or within 105m of the cave footprint (USFWS 2012).
- 12. Proximity to infrastructure/ utilities:
 - a. Subjective determination using aerial map that illustrates the cave's location and infrastructure within and adjacent to delineated preserve area boundaries.
 - b. Ideally preferred protection goal: cave preserve is free of underground pipelines, storage tanks, water retention ponds, or other structures/facilities that could cause contamination (USFWS 2012).
- B. <u>Hydrogeologic quality of troglobitic habitat measures</u>:
 - 1. Contribution to water quality/quantity within the karst ecosystem. Quantified by:
 - a. Surface Catchment Area size
 - b. Maximum Potential Subsurface Catchment Area based on data collected
 - c. Average combined drip rate per cave following methodology described in Hauwert and Cowan (2013).
 - d. Lack of subsurface pipelines or retention basins (USFWS 2012).
 - e. Ideally preferred protection goals: larger catchment areas are preferred due to their more significant contribution to water quantity within the karst ecosystem.
 - 2. Total accessed length, depth, and volume of cave:
 - a. Determined by cave maps. Also, if applicable, describe potential of undiscovered cave passages with supporting evidence.
 - b. Volume of cave determined by methods described in Krejca and Weckerly (2007).
 - c. Ideally preferred protection goal: USFWS (2012) states that larger, deeper caves may help protect against impacts to protected species from climate change by better maintaining in-cave stable temperatures and high humidity.
 - 3. Presence of permanent water bodies within cave:
 - a. Determined by in-cave surveys and/or documentation on cave maps and data from access to phreatic zone habitat where aquatic life such as aquatic salamanders may potentially be found.
 - b. Ideally preferred protection goal: caves with permanent bodies of water (pools, cave streams) are preferred for their contribution of habitat for aquatic life, potentially increasing biological diversity.

C. Ecological health measures:

These parameters may not be a critical factor on their own, but are important for the evaluation team to help understand the current ecological status and potential future management needs of the cave.

- 1. Healthy/stable cave cricket population:
 - a. Use existing cave cricket exit count survey results to assess population trends at caves.
 - b. If cave cricket data are absent or lacking, perform cave cricket monitoring following recommendations in USFWS 2014a.
 - c. Ideally preferred protection goal: results at caves will demonstrate a healthy and stable cave cricket population as demonstrated by repeated surveys.
- 2. Density of red-imported fire ants (RIFA), tawny crazy ants, and/or other invasive species that could impact the cave ecosystem:
 - a. Perform surveys using a scientifically accepted protocol for tawny crazy ants (*Nylanderia fulva*) to confirm absence at sites: caves suggested as candidates for replacement caves should not have infestations of tawny crazy ants.
 - b. Quantify RIFA densities using survey methods detailed in USFWS 2014a to ensure that RIFA threshold levels have not been reached at replacement caves.

Evaluation Documentation Requirements

The following documents and information should be included for conducting the evaluation to determine substitution need of a BCCP-listed cave and suitability of its replacement cave(s) (see above for details/definitions of specific factors):

- 1. Maps of each cave preserve area demonstrating the following:
 - a. Cave location and footprint.
 - b. Cave protection area.
 - c. Surface and subsurface drainage basin delineations.
 - d. Cave cricket 105 meter foraging area delineation.
 - e. Adjacent karst feature locations.
 - f. Landscape mosaic of karst preserve.
 - g. Incompatible forms of land use within the delineated karst preserve.
 - h. Infrastructure within and adjacent to delineated preserve area boundaries.
 - i. Surrounding land use.
- 2. Cave map for each cave demonstrating length, depth, and permanent bodies of water.
- 3. Documentation confirming presence of federally-listed karst invertebrates and/or BCCP karst SOC's.
- 4. Species lists for each cave.
- 5. In-cave faunal survey results demonstrating species abundance and methodology for conducting surveys.
- 6. Hydrogeologic study reports demonstrating methodology to assess drainage basin delineations and average combined drip rate results.
- 7. Cave cricket exit count survey results at caves and methodology for conducting surveys.
- 8. RIFA survey data for caves demonstrating mound densities.

See the Cave Comparison Worksheet below (Table 1) for a summary of cave substitution evaluation criteria.

Table 1 Cave Comparison Wo	orksheet
----------------------------	----------

Ca	ve Substitution Evaluation Criteria Worksheet	BCCP Cave	Substitution Cave	Comments
Sig	gnificant Diversity Criteria			
1a. Co	onfirmed endangered species locality?			
1a. Lis	st of endangered species (ES) present		and the second second	
1b. BC	CCP Species of Concern (SOC) locality?			Difference of the second of
1b. Lis	st of BCCP SOC present		tani i criates	Network - Calendial and Annako
a - 2c. Re	eplacement cave has similar or greater overall species diversity			
2d. Lis	st of additional troglobitic species			יעפר איבין אן 2005 איפרא אפעי
2d. SG	SCN list ranking of additional troglobitic species		office of special particular	neutropolitica de la construcción d
2d. Na	atureserve rarity rank of additional troglobitic species			
2e. Lis	st of rare non-troglobitic species			
3. Re	eplacement cave has similar or greater overall species abundance			C. P. D. M. Dir M. M. P.
	irst Fauna Region			12 St M20 Responsibility
	eplacement cave is within same BCCP cave cluster (if applicable)		Maint Mark	The show of the second
En	vironmental Protection Criteria			sphalests G. J. Luca
A. Ka	irst feature surface area protection measures:			A DOS TE LE STALL PROF
1. Pe	rcent of cave footprint within protected area			wind put and she calls
2. Dis	stance of cave footprint to nearest preserve edge			
3. Pe	ercent of surface drainage within protected area			
4. Pe	ccent of subsurface drainage within protected area		WITTER WITTER	
5. Pe	crcent of cave cricket foraging area within protected area			
6. Pre	eserve tract size			Without attended to a state
7. Ne	et gain in protected land for BCP			HI BOR ENGLAND
8. Shi	ape of protected area			states and states and services
9. Lai	ndscape mosaic of protected area (% woodland/grassland)	`		
10. Nu	umber of adjacent karst features within protection area	Contraction of the second		And the second second second second
	compatible land use/fragmentation			the Dictor and Stillers Dave
12. Pro	oximity to infrastructure/pipelines/utilities			and a satisfication of the state
B. Hy	drogeologic quality of troglobitic habitat measures:			
1a. Su	rface catchment area size			anna ann tulta an Antain
1b. Ma	aximum potential subsurface catchment area			is then an a transition Schema
1c. Av	verage combined drip rate	23	11. 11850 AS	PRODUCT PRODUCTION OF
2. To	tal accessed length of cave			
2. To	tal accessed depth of cave			RANGE STILLEN BRE GREAT
2. To	tal accessed volume of cave		and a sufficience	
3. Pre	esence of permanent water bodies in cave			
C Fre	ological health measures:			
	ve cricket population: in-cave survey and exit count results			
	ed-imported fire ant density at site wny crazy ants present at site?			

Literature Cited

Krejca, J.K. and F.W. Weckerly. 2007. Detection probabilities of karst invertebrates. Report prepared for Texas Parks and Wildlife Department and U.S. Fish and Wildlife Service.

Hauwert, NM, ME Litvak, and JM Sharp. 2005. Characterization and Water Balance of Internal Drainage Sinkholes: In Beck, B. ed., Geotechnical Special Publication No. 144, Sinkholes and the Engineering and Environmental Impacts of Karst, Proceedings of the Ninth Multidisciplinary Conference, p. 188-200.

http://www.austintexas.gov/watershed_protection/publications/document.cfm?id=186248

Hauwert, Nico M. 2009. Groundwater Flow and Recharge within the Barton Springs Segment of the Edwards Aquifer, Southern Travis County and Northern Hays County, Texas: Ph.D. Diss., University of Texas at Austin, Texas. 328 p. http://repositories.lib.utexas.edu/handle/2152/14107

Hauwert, N., and Cowan, B. 2013. Delineating Source Areas to Cave Drips and Cave Streams in Austin Texas, USA: 13th Sinkhole Conference, Carlsbad, NM. http://www.karstportal.org/node/11735?destination=node/11735

Master, L.L., D. Faber-Langendoen, R. Bittman, G.A. Hammerson, B. Heidel, L. Ramsay, K. Snow, A. Teucher, and A. Tomaino. 2012. NatureServe Conservation Status Assessments: Factors for Evaluating Extinction Risk. NatureServe, Arlington, VA.

Schneider, K. and D. C. Culver. 2004. Estimating subterranean species richness using intensive sampling and rarefaction curves in a high density cave region in West Virginia. Journal of Cave and Karst Studies 66(2): 39-45.

Taylor, S.J., J. Krejca, and M.L. Denight. 2005. Foraging range and habitat use of *Ceuthophilus secretus* (Orthoptera: Rhaphidophoridae), a key trogloxene in central Texas cave communities. American Midland Naturalist 154: 97-114.

Texas Memorial Museum (TMM). 2009. TEXBIO database, The University of Texas at Austin.

Texas Parks and Wildlife Department. 2011. Texas Conservation Action Plan: Species of Greatest Conservation Need. Austin, Texas. <u>https://tpwd.texas.gov/landwater/land/tcap/sgcn.phtml</u>

U. S. Fish and Wildlife Service. 1996a. Federal Fish and Wildlife Permit No. TE-788841-2. U. S. Fish and Wildlife Service. 2012. Karst Preserve Design Recommendations. Austin Ecological Services Field Office. Austin, Texas.

U.S. Fish and Wildlife Service. 2014a. Karst Preserve Management and Monitoring Recommendations. Austin Ecological Services Field Office. Austin, Texas.

U.S. Fish and Wildlife Service. 2014b. Section 10(a)(1)(A) Karst Invertebrate Survey Requirements for Conducting Presence/Absence Surveys for Endangered Karst Invertebrates in Central Texas. Fish and Wildlife Service, 10711 Burnet Road, Suite 200, Austin, Texas 78758. May 8, 2014.

Veni, G. 1992. Geological controls on cave development and the distribution of cave fauna in

the Austin, Texas, region. Report prepared for U.S. Fish and Wildlife Service, Austin, Texas. George Veni and Associates, San Antonio, Texas. 77 pp.

Veni, G. 2003. Delineation of hydrogeologic areas and zones for the management and recovery of endangered karst invertebrate species in Bexar County, Texas. Report for U.S. Fish and Wildlife Service, Austin, Texas, prepared by George Veni and Associates, San Antonio, Texas. Dated 23 December 2002 with minor revisions submitted 12 April 2003.

Zara Environmental LLC (Zara). 2013. Study Design: Cave cricket foraging study from Stark's North mine to (proposed) North Walnut Creek trail, Austin, Travis County, Texas. Report prepared for: RPS, 4801 Southwest Parkway, Austin, Texas 78735. 8 pp.

Zara Environmental LLC (Zara). 2014. Cave cricket foraging study from Stark's North Mine to proposed North Walnut Creek trail, Austin, Travis County, Texas. Report prepared for: RPS, 4801 Southwest Parkway, Austin, Texas 78735. 27 pp.

How is a Cave Substitution Made?

- a. A BCCP Coordinating Committee Member makes a proposal to substitute for a cave listed in condition S1 or T1 in the BCCP Federal Permit
- b. BCP staff(s) assembles information required, as described in the Evaluation Documentation Requirements
- c. Refer the proposal to the BCCP Scientific Advisory Committee Karst Subcommittee for review, assessment, and recommendation to the Coordinating Committee for action.
- d. When the Coordinating Committee takes action to accept a cave substitution proposal, the coordinating committee will initiate the BCCP amendment process for a minor amendment to the BCCP federal permit (Article 7, section 7.2, Interlocal Cooperation Agreement Between Travis County and City of Austin Implementing the Balcones Canyonlands Conservation Plan – Shared Vision)
- e. Proposals for cave substitution will be completed within one year from the date of submission by a Coordinating Committee member. This allows for any needed information gathering, additional field investigations, data analysis, and limited Coordinating Committee meeting schedules. This timeline will allow BCCP to appropriately address issues of non-compliance in a manner that would not result in an immediate permit violation while allowing third party actions to proceed in a reasonable manner.

Karst Preserve Protection and Management Measures

These management measures must be able to be enacted at BCCP caves or candidate replacement caves. If for some reason the candidate replacement cave's site can not adhere to these measures, then it may not be considered as a substitution.

- 1. No public access allowed in cave:
 - a. Ideally preferred protection goal: USFWS 2012 states that no public access should be allowed at caves: "to protect the subsurface habitat, several things should be carefully controlled including ensuring that the cave is entered for monitoring purposes only".
 - b. Candidate replacement caves should not allow public access.

- c. If recreational use is allowed in the cave's protected area, it should not interfere with karst management objectives as described in 11. c. (Incompatible land use) above and as defined in the most recently approved BCP Land Management Plan (Chapter IX, Karst Species Management).
- 2. Cave is (or will be) gated and/or fenced:
 - a. Ideally preferred protection goal: perimeter fencing around cave preserves is preferred for protection of the karst ecosystem from dumping, vandalism, and trespass (USFWS 2014a). Properly designed and installed cave gates are also preferred where there is a history of trespass and vandalism, and where human health or safety may be at risk (USFWS 2014a).
- Cave is (or will be) monitored/managed per most recent USFWS Karst Preserve Management and Monitoring Recommendations (2014a). Ideally preferred protection goals for this factor:
 - a. Biological monitoring is being conducted.
 - b. Vegetation management supports health of karst habitat.
 - c. Red-imported fire ant management is performed.



Record of Decisions

Balcones Canyonlands Conservation Plan Scientific Advisory Committee September 23, 2015, 3:00 PM to 5:00 PM Waller Creek Center, Room 104 625 East 10th Street Austin, Texas 78701

Present

Absent:

Jean Krejca

Sam Whitehead John Cornelius Craig Farquhar Nico Hauwert Sherri Kuhl, BCP Secretary Amie Treuer-Kuehn David Steed

John Cornelius, Vice Chair, led the meeting in Jean Krejca's absence.

- 1. **Approve Record of Decisions from regular meeting on 7/29/15**. Motion from Hauwert to approve, second by Steed, carried 6/0
- 2. **BCCP Cave Substitution Policy Update** Kuhl advised members that the Cave Substitution Policy was approved by the Coordinating Committee at their meeting on August 21, 2015. She discussed the steps that would be involved in the process and thanked everyone, especially the SAC karst subcommittee, for their time and efforts put into producing the process document. Nico stressed the importance of spending time and resources on looking for new caves and digging out features that have been previously filled in by ranchers.
- 3. Discuss and take action regarding a request from Coordinating Committee Member Pool to provide advice to the Coordinating Committee about whether or how SH45 SW might fragment habitat for wildlife in the vicinity of its proposed route - Krejca continues to work on providing a broad overview based upon available information from the FEIS and supporting documents.
- 4. Discuss and take action regarding a request from Coordinating Committee Member Pool to review surveys and other monitoring work for GCWA along SH45 SW in order to determine whether the effect call is supported by the data presented - Farquhar is reviewing Golden-cheeked Warbler information from the FEIS and supporting documents. Farquhar discussed the site visit made with City staff to the Water Quality Protection Land tract that the proposed SH 45 SW would cross. Hauwert updated the members on the subsurface drainage studies of Flint Ridge Cave, and the members discussed the potential SH 45 SW road project.

5. **Nominations for Chair and Vice-Chair for 2016** - Kuhl asked for nominations and selfnominations for Chair and Vice-Chair for 2016. Anyone interested in serving, please contact Kuhl and she will develop a slate of candidates for the first meeting of 2016.

6. Subcommittee reports:

- a. **Karst** Hauwert advised members that the City of Austin has reached agreement with the Village of Western Oaks community for the City to take over ownership and management of the Village of Western Oaks cave preserve; and are looking for a closing date. Watershed Protection Department will be the managing department and will continue using them for education. The repairs, restoration, and maintenance at Blowing Sink Preserve are ongoing, but almost complete. O'Donnell gave an update on the Tawny Crazy Ant Section 6 grant and upcoming research study on this new invasive species.
- b. **BCP Strategic Research Plan** Whitehead gave a report for the subcommittee and led a discussion on how to best move forward with developing a BCP Strategic Research Plan. First step will be to conduct a bibliography of all of the research that has been done on the BCP.

7. Staffs' reports:

- a. Lisa O'Donnell reviewed the written reports for the City and the BCP Partners Education and Outreach.
- b. Linda Laack and Melinda Mallia reviewed Travis County's written report.
- c. Erik Huebner reviewed LCRA's written report.
- 8. **Members' issues and concerns** Members discussed working on the State of the BCP report with the CAC. Whitehead and Treuer-Kuehn volunteered to help. It was also mentioned that there will be a Golden-cheeked Warbler and Black-capped Vireo Symposium in January of 2016.
- 9. Next SAC Meeting Kuhl discussed scheduling next year's meetings in January 2016.



Record of Decisions

Balcones Canyonlands Conservation Plan Citizens Advisory Committee Regular Meeting September 22, 2015 6:30 PM to 8:30 PM Waller Creek Center, Room 104 625 East 10th Street Austin, Texas 78701

Present
Peter Torgrimson, Chair
Forrest Arnold
Sherri Kuhl, BCCP Secretary
John Gosdin
Tom Hegemier
Annie Kellough

Richard DePalma Mary Ann Neely Kent Browning

2 Vacant

Excused Ken Beck David Whatley, Vice Chair

After call to order, Chair Torgrimson introduced Kent Browning as the new member appointed by Travis County, and Richard DePalma from the City of Austin Parks and Recreation Board. All the members introduced themselves, and Sherri Kuhl described the process in place to confirm the City of Austin Boards and Commission Members by the Austin City Council.

1. Citizens Communication None

2. BCP Partner Reports

- a. **City of Austin BCP** Lisa O'Donnell reviewed the City's written report with the committee and the BCP Partners Education and Outreach Report.
- b. **Travis County BCP** Melinda Mallia and Linda Laack reviewed the Travis County written report with the committee.
- c. LCRA BCP Erik Huebner reviewed the LCRA written report with the committee.
- d. **Other BCP Partners** Forrest Arnold gave an update on the Violet Crown Trail for the City of Sunset Valley.

3. New Business

a. **Members' issues and concerns** – Members requested that we do a briefing for new members on the BCCP. Sherri Kuhl said that BCP staff will work on this in the new year. There was discussion about the time allotted for speakers, the meeting location, and meeting member attendance. A question was raised about whether these meetings were subject to the Open Meetings Act and the Public Information Act. Sherri confirmed that they are not subject to the Open Meetings Act. The agenda and meeting information are posted as a courtesy to the public. Kuhl agreed to look into the issue of whether communications between members are subject to the Public Information Act and to report back.

4. Old Business

- a. **Approve the Record of Decision from 7/28/15** Motion to approve from Gosdin, Second Hegemier, Carried 5/0, with the new members from the City Boards and Commissions abstaining from voting until they are approved by City Council.
- b. CAC action regarding positive press coverage from 10/22/14 Kuhl noted that staff is working with PIO and others to advance this recommendation. Distributing a large BCCP article to the Community Impact papers and/or Four Points News will be timed with the 20th Anniversary in May 2016.
- c. Funding concerns for rural and small community fire departments adjacent to BCP Mallia reported on the July 30, 2015 meeting of the task group formed from the Austin Travis County Joint Wildfire Coalition.
- d. Update on BCCP Karst Substitution Process planning and take action as appropriate Kuhl advised members that the Cave Substitution Policy was approved by the Coordinating Committee during their August 21, 2015 meeting.
- e. CAC vacant positions and expected changes from City Boards and Commission changes - Kuhl advised members that two consensus positions are vacant and she will be convening the Advisory Committee Task Group soon to develop recommendations for the two CAC consensus positions.
- f. **CAC Annual Work Plan -** John Gosdin recommended that work be focused on the State of the BCP report this year and to remove the other two items from the CAC Annual Work Plan. Peter Torgrimson reported on the purpose of the subcommittees and in discussion it was agreed to focus on the State of the BCP report and form a subcommittee with the SAC and staff to work on it, with a completion date of May 2, 2016.
 - i. Report from subcommittees:
 - Wildfire and fire support.
 - Invasive species in landscaping.
 - State of BCP report (joint with SAC).

City of Austin BCP Report BCCP Coordinating Committee Meeting August 1, 2015 – September 15, 2015

Program Administration

Acquisition

• The 10-acre Lucas tract was purchased for addition to the City BCP.

Personnel

• Willy Conrad retired from his position as Division Manager of the Wildland Conservation Division on August 31, 2015. Sherri Kuhl is acting BCCP Coordinating Committee Secretary and Kevin Thuesen is acting Wildlands Division Manager. Two positions will be filled and the BCCP Coordinating Committee Secretary will be jointly funded by the City of Austin and Travis County.

BCCP Infrastructure Training Workshop

• Scheduled for October 8, 2015, at the Lady Bird Johnson Wildflower Center.

Biological Monitoring

Karst

- Tawny crazy ants (TCAs) were first observed negatively impacting cave fauna in Whirlpool Cave in June 2013, and on November 20, 2014, TCA's were discovered in No Rent Cave. In an effort to monitor impacts, City and Travis County staff continues to monitor Whirlpool Cave as well as No Rent and Weldon Cave (in close proximity to TCA's). We still anticipate that this new invader will have a major negative effect on cave fauna. The section 6 grant proposal (investigate possible methods of treating TCAs within the cave cricket foraging area) was awarded, with a start date of October 2015.
- BCCP Cave Substitution Policy was officially adopted by the BCCP Coordinating Committee on August 21, 2015. This new policy provides the groundwork for possible future substitutions for BCCP permit caves.
- Conducted cave faunal surveys and cave cricket exit counts (ongoing).
- Monitored cave conditions, removing trash, and treating for red-imported fire ants (ongoing).
- BCP Staff is continuing to assist Nico Hauwert and the Watershed Protection Department with the Blowing Sink critical environmental feature (CEF) stabilization CIP project. Several major recharge features/caves are unstable, leading to large amounts of sediment washing into the aquifer. The project will stabilize these features, thus improving water quality to Barton Springs. This project is nearing completion.
- In an effort to educate the public about the importance of caves and cave fauna, Nico Hauwert initiated an effort to find local non-BCP caves that would be appropriate for educational purposes. Staff and volunteers are currently excavating several karst features. Several significant caves have been opened including grassy cove cave which was recently gated. (ongoing)
- Cave public access: A cave collaboration group of City and County staff and volunteers met regarding guidelines for future public access to local caves including some BCP permit caves. The goal is to try to determine which caves are appropriate for public access, determine the appropriate number allowed per cave, and come up with a permitting system that would determine qualifications needed for leading cave tours (ongoing).
- Restoration and monitoring efforts have begun at 2 endangered species caves located in the courtyard at McNeil High. Nico Hauwert and McNeil High School science teacher Tina Vick are leading the restoration effort along with help from FWS, City and Travis County BCP staff. Travis County and City BCP staffs are currently conducting cave faunal surveys and cricket exit counts with the goal of including student participation for long term monitoring and potential research projects that will benefit the listed species. Travis County and City BCP staff has also held 2

volunteer clean up dates, removing large volumes of trash. City staff has provided plants for this project, planting 86 trees and shrubs, many of which provide forage for cave crickets (ongoing).

Golden-Cheeked Warbler (GCWA)

- <u>USFS Project</u>: Interlocal Agreement with the US Forest Service (Dr. Frank Thompson, Jennifer Reidy) to provide GCWA Population Viability and Habitat Suitability modeling within the BCP.
 - 5-year study, focuses on four primary questions:
 - What is the absolute abundance of the GCWA on the BCP and on individual macrosites?
 - How do demographics (e.g. density, productivity, survival) vary with landscape and habitat factors?
 - How viable are these populations?
 - How do various management scenarios influence population viability?
 - Fifth year of data have been sent to USFS; work on the models is in progress.
- The following 3 publications are now available online:
 - Reidy, J., F. Thompson, C. Amundson, and L. O'Donnell. 2015. Landscape and local effects on occupancy and densities of an endangered wood-warbler in an urbanizing landscape. Landscape Ecology; DOI: 10.1007/s10980-015-0250-0.
 - Reidy, J., L. O'Donnell, and F. Thompson. 2015. Evaluation of a reproductive index for estimating songbird productivity: case study of the Golden-cheeked Warbler. Wildlife Society Bulletin; DOI: 10.1002/wsb.576.
 - O'Donnell, L., C. Farquhar, J. Hunt, K. Nesvacil, J. Reidy, W. Reiner, J. Scalise, and C. Warren. 2015. Density influences accuracy of model-based estimates for a forest songbird. Journal of Field Ornithology; DOI: 10.1111/jofo.12116.

Black-capped Vireo (BCVI)

• COA BCP staff and volunteers are continuing to implement demonstration projects at the Vireo Preserve to promote slope stabilization, soil restoration, and regeneration of native plants on degraded areas that are currently not habitat for either the GCWA or BCVI. Since Vireo Preserve supports many of the habitat types observed throughout the BCP, lessons learned from restoration work at this site should be applicable to other areas within the BCP.

Jollyville Plateau Salamander (JPS)

- COA BCP staff assisted the Watershed Protection Department with several JPS surveys; most surveys were postponed due to recent floods. (ongoing)
- BCP staff conducted a salamander survey in Testudo Tube, the survey area is a 100 meter transect, all salamanders were captured, measured and photographed with the goal of identifying individuals via this set of photos since each individual salamander has distinct markings on the dorsal side of head. The ability to identify individual salamanders will allow for a better understanding of the salamanders life history and future trends.

Bracted Twistflower

• Plans for fence to protect Bracted Twistflower at Mount Bonnell: see Dual-Managed Lands, below.

Preserve Management

Dual-Managed Lands

• AWU and PARD are working to close and restore trails within the Connors Creek critical water quality protection zone in Emma Long Motocross Park. Additional signs are scheduled for installation. Staff is developing a stream assessment, restoration and monitoring plan with assistance from staff in the Watershed Protection Department.

- The proposed Mount Bonnell fence to protect bracted twistflower habitat and keep people off the bluff has been designed and construction will begin in the fall.
- PARD and BCP staffs worked with the Hill Country Conservancy on the installation of 911/mile markers in the Barton Creek Greenbelt.
- Barton Creek homeless camps: see Law Enforcement below.

Infrastructure Management

- LCRA successfully completed their upgrade of the T-160 line which runs through the Bull Creek macrosite and received construction approval from BCCP. LCRA worked closely with BCP and WPD staff to avoid impacts to endangered/threatened species and habitats. LCRA staff is currently addressing some erosion issues with guidance from BCP staff.
- BCP is currently working with the West Travis County PUA (WTCPUA) on construction of a new water service line which includes drilling underneath Barton Creek. (ongoing)

Law Enforcement

• Homeless camps are an ongoing management challenge in the Barton Creek Greenbelt. City Wildland Conservation and PARD staffs continue to monitor these sites and remove camps as they are continue to break into the Lanier house and that structure is scheduled for demolition this fall.

Invasive Species/Animal Management

- See Tawny Crazy Ants, above.
- BCP staff is continuing efforts to remove non-native plants on COA BCP tracts.
- BCP staff is actively removing feral hogs.

Oak Wilt

• BCP staff continued to monitor movement of the fungus in known oak wilt centers and success of control trenches.

Fencing Projects

- Bids for fence installation on the Jester and Cortaña tracts have been received and work is close to completion on the Jester game fence.
- Currently scoping surveys to fence the newly acquired Lucas tract, a portion of the Forest Ridge tract, Long Canyon, and Park West.

Fire Management

- Wildland Conservation Division (WCD) staffs have created a final draft of Wildfire Contingency Plan Maps for all WCD properties. These maps are intended to facilitate wildfire suppression on WCD property by illustrating access points, roads, structures, vegetation, and fire potential.
- WCD staff is currently planning additional potential shaded fuel break projects for the fall and winter.

	BCCP Administration			
Public Participation	Ten BCCP applications were received and processed; Two Participation Certificates were issued for a total of \$4,750.00 in mitigation payments. Fees were split 50:50 between Travis County and City of Austin.			
Land Acquisition	The 500-acre Kotrla tract in northwest Travis County was acquired in July. This tract is connected to another 450-acre BCP tract and this combined area provides complete protection of the surface and subsurface drainage basins of the BCCP-listed Spanish Wells cave. This area also contains habitat for warblers, vireos, and a population of Texabama croton, a plant named for protection in the BCCP permit.			
	Meetings have been held with the City of Jonestown to discuss City Council's concerns that the purchase of the 228-acre Plains Capital Tract for the BCP has negatively impacted anticipated revenue from development of a residential subdivision on that tract. The tract was purchased February 25, 2015.			
Personnel	Four Natural Resources positions were filled during August and September. David Morier is the new Project Manager; Julie Murray and Blake Sissel are Natural Resources Specialists; and Travis Clark is the new Natural Resources Tech. Hiring is in the final stages for the vacant Environmental Specialist position. A Community Liaison position will be advertised in October.			
	Rare and Endangered Resources Management			
Golden-cheeked warbler and Black-capped vireo	Warblers and vireos have migrated south for the winter. Nesting season officially ended August 31 st .			
Karst invertebrates	Summer season faunal surveys were performed at nine caves.			
	Summer season cave cricket exit counts were performed at six caves.			
	BCCP Cave Substitution Policy was adopted by the BCCP Coordinating Committee in August. This policy provides a process that allows caves listed in the BCCP permit to be substituted with other suitable caves in a manner that is transparent, science based, and consistent with the vision and intent established for BCCP.			
Plant and Animal Control				
Feral hogs	Signs of hog damage were found on many BCP tracts. Traps were in operation on the Ribelin, Sam Hamilton, and Concordia tracts.			
White-tailed deer	Spotlight deer surveys are scheduled for September and October. Deer harvest will begin in October.			
Tawny crazy ants	Assessments and surveys of Tawny crazy ant infestation were conducted at Whirlpool and No Rent caves.			

Invasive Plants	Invasive plants (<i>Nandina,</i> Brazilian Vervain, and Asian jasmine) were removed from the Snowden tract.		
Land Stewardship			
Fencing	Fencing projects are planned for this fall/winter on the Collins, Plains Capital, Scott and White, and Kotrla tracts.		
Wildfire Preparedness	Fuel mitigation projects are planned for this fall/winter on the Steiner Ranch, Canyon Vista, Greenshores, Grandview Hills, and Concordia tracts.		
	FEMA has approved the final award for a Hazard Mitigation Grant Program (HMGP) project to mitigate wildfire in the Wildland Urban Interface (WUI) of the BCP using shaded fuel treatments. The grant project will begin upon receipt of an award notice from the Texas Division of Emergency Management.		
	With or without the grant, plans are underway to begin several new wildfire mitigation projects in the BCP WUI after October 1, using funds approved by Commissioners Court in the Natural Resources Program budget.		
Infrastructure Corridor	LCRA conducted inspections and maintenance of the T-160 and T-161 transmission lines which run through the Steiner, Ribelin, and Lucas Tracts.		
	An Infrastructure Workshop is scheduled for October 8 th at the Wildflower Center. This workshop is designed to help explain the significance of the BCP and provide guidance to infrastructure service providers that cross the BCP or that pass through endangered species habitat outside of the preserve.		
Transportation	The status of the following transportation projects have not changed since the last staff report:		
	LISD is pursuing an individual ESA 10(A) permit for the proposed road project in the infrastructure corridor through Travis County Ribelin, Sam Hamilton East, and private 10(A) mitigation lands.		
	Some traffic improvements have been made along RM 2222. Additional improvements involving RM2222 and RM 620 are still in development. There has been no recent discussion of the LISD's proposed new roadway.		

	SH 45 SW: Final Environmental Impact Statement (FEIS) was released January 23, 2015. The Record of Decision dated March 4 th identified the
	Preferred Alternative as the Selected Alternative. See http://www.sh45sw.com/about/environment.php
	USFWS responded to letters from City and County elected officials re concerns. The future status of Flint Ridge Cave remains in question. Details of land management in the vicinity of FRC remain to be resolved. On April 21 st the Travis County Commissioners Court directed staff to work with the Central Texas Regional Mobility Authority (CTRMA) to include state-of-the-art stormwater controls for this area of extraordinary sensitivity; and to work with the CTRMA to require a sharing, by the CTRMA, of the risk of non-compliance under the Endangered Species Act.
	TNR staff are currently engaged in ongoing review of project plans. TNR staff are also participating in a number of technical working groups with CTRMA, City staff and other stakeholders to review and assess design plans, water quality controls, protection of Flint Ridge Cave and other related issues.
	Travis County commissioned a feasibility study for the realignment of Steiner Ranch Boulevard/Comanche Trail. The recommendation rejected any new encroachments on BCP and proposed potential new alignments, but these were contingent on the results of the TxDOT RM 620 Corridor Study which is still in progress.
Adjacent Development	Staff coordinated with City of Austin and developers on current and future development projects adjacent to the Cuevas, Nootsie, The Crossings, and Steiner Ranch tracts. Staff is reviewing fire service road route options for the Travaasa development. The road will cross through The Crossings tract.
	Law Enforcement
Trespass and Criminal Mischief	Four incidents of construction material dumping was discovered on the Cuevas and Plains Capital tracts; Vandalism (damage to PVC pipes) occurred on the Webb tract.
	A 0.3 acre area of tree clearing on the Gray Mountain conservation easement was observed in May. After negotiations, the owner has agreed to restore vegetation on the cleared area.
	Outreach/Volunteers
Volunteers provided September.	80.5 hours of service to the Travis County BCP during August and

Lower Colorado River Authority (LCRA) Balcones Canyonlands Preserve (BCP) Update for McGregor, Wheless and Westcave Preserves

BCCP CAC/SAC Meetings – August 22-23, 2015

Wildlife Surveys at Wheless

LCRA worked with an outside contractor to perform wildlife surveys to determine white-tailed deer population density, buck/doe ratio and fawn survival. A final report is in progress, which will include findings, visibility measurements and copies of completed data sheets for each survey attempt. Incidental observations of other species will be noted in the report.

Erosion Control Work on City of Austin (COA) 3M Preserve for LCRA T-160 Project

LCRA rebuilt the T-160 (Marshall Ford to McNeil) transmission line, which crosses several BCP lands. Concerns arose over erosion control on the 3M preserve that resulted from the construction of the transmission line. LCRA worked with City of Austin (COA) BCCP staff to finalize and implement an erosion control plan for this preserve. Implementation of the erosion control plan began in September 2015 and will continue until early October 2015. Some of the implemented erosion controls include gabion mattresses and berms where pad sites construction occurred (see photos below). Top soil from cave restoration work was removed from various locations in southwest Austin and will be used with a BCCP-approved seed mix from Native American Seed to help stabilize the pad sites. Road work will be conducted and multiple diversion berms created to assist in slowing sheet flow from rain events.





Westcave Preserve Land Management Activities

American Youth Works cut understory woody species in the future prairie area, in preparation for a USFWS prescribed burn. They also established prairie restoration area photo points.

Campers with El Ranchito Service Corps erected a bat box, removed Johnsongrass, and performed brush management on the uplands. The group also cut and treated Chaste Tree (*Vitex agnus-castus*) directly above the waterfall in the upper creek.

Girl Scouts researched and developed laminated informational sheets on birds that can be seen in the area and built a wooden stand to hold the information sheets.

A dragonfly research study was conducted through Migratory Dragonfly Partnership for Citizen Scientists and has been set up in two places on the preserve and one at Hammett's Crossing. Westcave conservation staff will monitor monthly.

Salamander Survey conducted at Westcave with TPWD, Travis County and COA biologists. Biologists found no salamanders on the preserve, but found a unique species at Reimers Ranch. Staff conducted an additional survey but no salamanders were observed.

General trail maintenance and boundary inspections were conducted as well as an endemic plant survey. Westcave staff is preparing for the avian population monitoring project to be conducted by Dawn Houston, a member of the Conservation Committee. The project includes conducting avian point counts throughout the Preserve to monitor population and trends.

Illegal Dumping

An LCRA Transmission Line crew was inspecting the T-142 (Hi Cross to Marshall Ford) transmission line. This line crosses several BCPs. The crew discovered illegal dumping in the transmission line easement at the edge of the Barton Creek Park. LCRA reported the dumping to COA BCCP biologist who informed LCRA that the information was given to staff in the COA Parks and Recreation Department, as they are technically the owners at the location where the dumping occurred.



Exotic Plants on Wheless

Update: COA biologists plan to meet with LCRA between October and December 2015 to flag and discuss the management of exotic plant species.

In June 2015, COA Endangered Species Biologist completed a helicopter survey of the BCP and informed LCRA that no oak wilt was observed on the Wheless Preserve, but they did find some Tree-of-Heaven (*Ailanthus altissima*). The biologist requested access to Wheless to flag the exotic trees or collect GPS points of their locations to aid LCRA staff when managing exotics. The biologist also requested to check the flow for the Jollyville Plateau Salamanders located on the preserve. LCRA granted access, but the biologist have not entered the preserve to date (6/18/15).



PARTNER EDUCATION SUMMARY August 1 - September 15, 2015



- City of Austin staff hosted a volunteer CPR/First Aid training for volunteers that lead projects independently on BCP properties.
- Staff is starting to plan a BCCP 20th Anniversary Event.
- Upcoming events include the BCCP Infrastructure Workshop 10/8 8am-12:30pm
 - http://2015bccpworkshop.eventbrite.com

Guided Hikes			
		#	
DATE	ACTIVITY	PARTICIPANTS	PARTNER
8/8/2015	2nd SAT. Guided Hikes	15	WBSN BCP
9/12/2015	2nd SAT. Guided Hikes	12	WBSN BCP
8/22/2015	TAS Bird Hike	16	TAS BCP
9/14/2015	Sun City Hiking Club	19	TAS BCP
8/8/2015	2nd SAT. Guided Hikes	15	WBSN BCP
9/12/2015	2nd SAT. Guided Hikes	12	WBSN BCP
8/22/2015	TAS Bird Hike	16	TAS BCP
2015 Summer	QTR. YOUTH EDU PARTICIPANT TOTAL:	105	
YEAR TO DA	ATE YOUTH EDU PARTICIPANT TOTAL:	3069	

BCP HIKE AND LECTURE SERIES				
DATE	ACTIVITY	# PARTICIPANTS	PARTNER	
8/1/2015	Flint Knapping	10	WBSN BCP	
8/5/2015	Night Hike at Sam Hamilton	10	COA BCP	
9/5/2015	Spider Joe	14	WBSN BCP	
2015 Summer QTR. HIKE AND LECTURE PARTICIPANT TOTAL: 34				
YEAR TO DATE HIKE AND LECTURE PARTICIPANT TOTAL: 199				

BCP gains valuable help completing tasks with the help of numerous volunteers. Below is a summary of volunteer activities from August 1st through August 15th. Tracking volunteer hours includes updates and additions after the printing of this report. The most up-to-date volunteer hours will be available in the year-end report.

Volunteer Summary				
Total Vol.				
Weekly volunteer stewardship group (sum for Aug)		WB BCP		
Collecting Veg plot data	42.5	TAS BCP		
Facilities Maintenance	4	TAS BCP		
Trail Maintenance	7	TAS BCP		
Grounds Maintenance	1.5	TAS BCP		
Habitat Restoration	13	TAS BCP		
Invasive Plant Removal	9	TAS BCP		
Travis County Volunteer Projects	80.5	TC BCP		
Land Stewardship Days at the Vireo Preserve	84	COA BCP		
Tree Mulching	8	COA BCP		
CPR and First Aid for Volunteers	36	COA BCP		
Saturday stewardship day (Aug)	60	WB BCP		
Pollinator Garden and Tree Mulching Project	164	COA BCP		
Weekly volunteer stewardship group (sum thru Sept 17)	6	WB BCP		
e Summer QTR. VOL ACTIVITIES TOTALS:	2,204			
	Weekly volunteer stewardship group (sum for Aug) Collecting Veg plot data Facilities Maintenance Trail Maintenance Grounds Maintenance Habitat Restoration Invasive Plant Removal Travis County Volunteer Projects Land Stewardship Days at the Vireo Preserve Tree Mulching CPR and First Aid for Volunteers Saturday stewardship day (Aug) Pollinator Garden and Tree Mulching Project Weekly volunteer stewardship group (sum thru Sept 17)	Total Vol. Hours Weekly volunteer stewardship group (sum for Aug) 6 Collecting Veg plot data 42.5 Facilities Maintenance 4 Trail Maintenance 7 Grounds Maintenance 1.5 Habitat Restoration 13 Invasive Plant Removal 9 Travis County Volunteer Projects 80.5 Land Stewardship Days at the Vireo Preserve 84 Tree Mulching 8 CPR and First Aid for Volunteers 36 Saturday stewardship day (Aug) 60 Pollinator Garden and Tree Mulching Project 164 Weekly volunteer stewardship group (sum thru Sept 17) 6		